

A GENERALIZATION OF THE BRAUER ALGEBRA

WILLIAM Y.C. CHEN AND CHRISTIAN M. REIDYS^{*}

ABSTRACT. We study two variations of the Brauer algebra $\mathbb{B}_n(x)$. The first is the algebra $\mathbb{A}_n(x)$, which generalizes the Brauer algebra by considering loops. The second is the algebra $\mathbb{L}_n(x)$, the $\mathbb{A}_n(x)$ -subalgebra generated by diagrams without horizontal arcs. $\mathbb{A}_n(x)$ and $\mathbb{L}_n(x)$ exhibit for $x \neq 0$ an hereditary-chain indexed by all integers. Following the ideas of Martin [10] in the context of the partition algebra, and Doran *et al.* [4] for the Brauer algebra, we study semisimplicity of $\mathbb{A}_n(x)$ using restriction and induction in $\mathbb{A}_n(x)$ and $\mathbb{L}_n(x)$. Our main result is that $\mathbb{A}_n(x)$ is semisimple if $x \notin \mathbb{Z}$ and that $\mathbb{L}_n(x)$ is semisimple if $x \neq 0$.

1. INTRODUCTION

In this paper, we study the semisimplicity of the two diagram algebras $\mathbb{A}_n(x)$ and $\mathbb{L}_n(x)$. $\mathbb{A}_n(x)$ generalizes the Brauer algebra, $\mathbb{B}_n(x)$, by containing diagrams in which vertices can be incident to loops (or equivalently, isolated vertices). $\mathbb{L}_n(x)$ is the $\mathbb{A}_n(x)$ -subalgebra generated by all diagrams without any horizontal arcs. The motivation for considering these algebras is twofold: on the one hand in the context of Schur-Weyl duality: $\mathbb{A}_n(x)$ is the centralizer algebra of the group of stochastic, orthogonal matrices and $\mathbb{L}_n(x)$ is the centralizer algebra of the group of stochastic, invertible matrices. On the other hand, $\mathbb{A}_n(x)$ is as the algebra of partial matchings of importance for RNA pseudoknot structures, i.e. helical configurations of RNA primary sequences with cross-serial nucleotide interactions [8].


The Brauer (centralizer) algebras [3] over the field $F = K(x)$, denoted by $\mathbb{B}_n(x)$, are finite dimensional F -algebras indexed by a positive integer n and x , which is either algebraic or transcendent over K . $\mathbb{B}_n(x)$ is the centralizer algebra for the orthogonal or symplectic group on the n th tensor powers of the natural representation. $\mathbb{B}_n(x)$ has been studied by various authors mainly using

Date: February, 2009.

combinatorial methods, see [1, 2, 13, 6, 5, 7] and [15]. Hanlon and Wales conjectured that $\mathbb{B}_n(x)$ is semisimple for all $x \notin \mathbb{Z}$ [6]. Their conjecture was proved by Wenzl [16] and Rui [11] gave necessary and sufficient conditions for the Brauer algebras to be semisimple.

The analysis presented here is based on the concepts of Martin [10] developed in the context of the partition algebra, \mathbb{P}_n . Martin's key idea was to relate the existence of certain embeddings to semisimplicity. Subsequently, Doran *et al.* [4] used this framework in order to offer an alternative to Wenzl's proof of semisimplicity. Wenzl's inductive construction hinges on an interpretation of a key ideal in $\mathbb{B}_n(x)$ as the tensor product $\mathbb{B}_{n-1}(x) \otimes_{\mathbb{B}_{n-2}(x)} \mathbb{B}_n(x)$ [9] and the nondegeneracy of a Markov-trace arising naturally in the construction of the latter. The nondegeneracy of this trace form is a result of Weyl's character formulas and is in this sense somewhat "unsatisfactory". The work of Martin [10] and Doran *et al.* [4] puts semisimplicity in the context of quasi-hereditariness and allows to avoid the use of Markov-traces.

Let \mathcal{A}_n be the set of partial 1-factors over $2n$ vertices, i.e. graphs over $2n$ vertices in which each vertex has either degree one or zero. We refer to \mathcal{A}_n -elements as diagrams and represent them by arranging the $2n$ vertices in two rows, each containing n vertices, with the rows arranged one on top of the other. Furthermore, we equip each isolated vertex with a loop. The n top-vertices are labeled by $[n] = \{1, \dots, n\}$ in increasing order and the n bottom-vertices are labeled by $[n'] = \{1', \dots, n'\}$. Let $\mathcal{L}_n \subset \mathcal{A}_n$ be the subset of all \mathcal{A}_n -diagrams without any horizontal arcs. We denote the subset of \mathcal{L}_n -diagrams having only vertical arcs by \mathcal{S}_n . When drawing diagrams, we oftentimes omit vertex labels. For instance,



are particular \mathcal{A}_n -, \mathcal{L}_n - and \mathcal{S}_n -diagrams. By abuse of notation, we write S_n instead of \mathcal{S}_n , identifying S_n with its embedding into \mathcal{A}_n . As for their cardinalities we immediately compute


$$(1.1) \quad |\mathcal{A}_n| = \sum_{j=0}^n \binom{2n}{2j} \prod_{i=0}^{(n-j)-1} (2(n-j) - 1 - 2i) \quad \text{and} \quad |\mathcal{L}_n| = \sum_{j=0}^n \binom{n}{j} \binom{n}{j} (n-j)!,$$

where the factor $\prod_{i=0}^{(n-j)-1} (2(n-j) - 1 - 2i)$ equals the dimension of the Brauer-algebra $\mathbb{B}_{n-j}(x)$. Arcs joining two different vertices, contained both in the top or bottom row are called horizontal arcs. Arcs joining top- and bottom-vertices are called vertical arcs. The induced subgraph of the top and bottom row of a diagram \mathbf{a} is denoted by $\text{top}(\mathbf{a})$ and $\text{bot}(\mathbf{a})$. Let \mathbf{e}_i be the diagram having

straight verticals except of the horizontal arcs connecting $i, i+1$ and $i', (i+1)'$, \mathbf{u}_i having straight verticals and loops at i and i' and \mathbf{g}_i having straight verticals except of the vertical arcs $(i, (i+1)')$ and $(i+1, i')$. Pictorially,

$$\begin{array}{c} \mathbf{e}_i = \begin{array}{ccccccc} 1 & \cdots & i & i+1 & \cdots & n \\ \bullet & & \bullet & \bullet & & \bullet \\ \vdots & & \vdots & \vdots & & \vdots \\ 1' & \cdots & i' & (i+1)' & \cdots & n' \end{array} \quad \mathbf{u}_i = \begin{array}{ccccccc} 1 & \cdots & i-1 & i & i+1 & \cdots & n \\ \bullet & & \bullet & \bullet & \bullet & & \bullet \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ 1' & \cdots & (i-1)' & i' & (i+1)' & \cdots & n' \end{array} \quad \mathbf{g}_i = \begin{array}{ccccccc} 1 & \cdots & i & i+1 & \cdots & n \\ \bullet & & \bullet & \bullet & & \bullet \\ \vdots & & \vdots & \vdots & & \vdots \\ 1' & \cdots & i' & (i+1)' & \cdots & n' \end{array} \end{array}$$

We now describe the multiplication of two diagrams. Let x be either a K -transcendent or algebraic element. We consider $F[\mathcal{A}_n]$, the free F -module generated by \mathcal{A}_n and show that $F[\mathcal{A}_n]$ is a monoid whose multiplication extends that of $\mathbb{B}_n(x)$ in a natural way. To this end let $\mathbf{a}, \mathbf{b} \in \mathcal{A}_n$. Let $G(\mathbf{a}, \mathbf{b})$ be the graph obtained by arranging the diagram \mathbf{a} above \mathbf{b} and introducing the vertical arcs (i, i') , $1 \leq i \leq n$ where i and i' are contained in $\text{top}(\mathbf{a})$ and $\text{bot}(\mathbf{b})$ -vertex, respectively. For instance,



$G(\mathbf{a}, \mathbf{b})$ contains two types of information: (i) $\ell(\mathbf{a}, \mathbf{b})$, the number of $G(\mathbf{a}, \mathbf{b})$ components that do not contain any vertices of $\text{top}(\mathbf{a})$ or $\text{bot}(\mathbf{b})$ and (ii) $G'(\mathbf{a}, \mathbf{b})$, the graph over the $\text{top}(\mathbf{a})$ and $\text{bot}(\mathbf{b})$ -vertices obtained as follows: any two vertices are connected by an arc if and only if they are connected by a $G(\mathbf{a}, \mathbf{b})$ -path. Accordingly, we have $\mathbf{a} \cdot \mathbf{b} = x^{\ell(\mathbf{a}, \mathbf{b})} G'(\mathbf{a}, \mathbf{b})$ and we shall write $\mathbf{a}\mathbf{b}$ instead of $\mathbf{a} \cdot \mathbf{b}$. $F[\mathcal{A}_n]$ becomes via “ \cdot ” an associative, unitary F -subalgebra of the partition algebra, which we denote by $\mathbb{A}_n(x)$. Furthermore, via “ \cdot ”, $F[\mathcal{L}_n]$ becomes an associative F -subalgebra of $\mathbb{A}_n(x)$, denoted by $\mathbb{L}_n(x)$.

By abuse of notation, we write $\mathbb{A}_n = \mathbb{A}_n(x)$, $\mathbb{B}_n = \mathbb{B}_n(x)$ and $\mathbb{L}_n = \mathbb{L}_n(x)$. Furthermore, we shall assume that F is a field of characteristic zero and the term “semisimple” is synonymous to “direct sum of full matrix algebras”. In other words, F is a splitting field of \mathbb{A}_n and \mathbb{L}_n .

Remark 1. Let $\ell_1(\mathbf{a}, \mathbf{b})$ and $\ell_2(\mathbf{a}, \mathbf{b})$ denote the number of inner components that are cycles and lines with loops at the start and endpoint. Setting

$$(1.2) \quad \mathbf{a} \circ \mathbf{b} = x_1^{\ell_1(\mathbf{a}, \mathbf{b})} x_2^{\ell_2(\mathbf{a}, \mathbf{b})} G'(\mathbf{a}, \mathbf{b}),$$

we observe that $F[\mathcal{A}_n]$ becomes via “ \circ ” an associative unitary F -algebra, which we denote by $\mathbb{A}_n(x_1, x_2)$. Obviously, in case of $x_1 = x_2$ the multiplications “ \circ ” and “ \cdot ” coincide.

As it is the case for \mathbb{B}_n , there exist natural embedding between \mathbb{A}_{n-1} and \mathbb{A}_n obtained by adding the vertices n and n' together with the straight vertical arc, (n, n') , $\epsilon_n: \mathbb{A}_{n-1} \longrightarrow \mathbb{A}_n$. By restriction the latter induces an embedding of \mathbb{L}_{n-1} into \mathbb{L}_n , which we denote again by $\epsilon_n: \mathbb{L}_{n-1} \longrightarrow \mathbb{L}_n$. Furthermore, there exists an involution on \mathbb{A}_n and \mathbb{L}_n obtained by transposing the rows, denoted by $\mathfrak{a} \mapsto \mathfrak{a}^*$. We set $\mathcal{A}_n^m \subset \mathcal{A}_n^n = \mathcal{A}_n$ to be the subset of diagrams having at most m vertical arcs and let \mathbb{A}_n^m be the ideal generated by \mathcal{A}_n^m . The ideals \mathbb{A}_n^m for $0 \leq m \leq n$ give a filtration of \mathbb{A}_n , i.e. we have

$$(1.3) \quad \mathbb{A}_n^0 \subsetneq \mathbb{A}_n^1 \subsetneq \cdots \subsetneq \mathbb{A}_n^{n-1} \subsetneq \mathbb{A}_n^n = \mathbb{A}_n.$$

Furthermore, let $\mathbb{I}_n^m = \mathbb{A}_n^m / \mathbb{A}_n^{m-1}$ denote the algebra induced by \mathbb{A}_n , which is generated by the set all \mathcal{A}_n -diagrams with exactly m vertical arcs, denoted by \mathcal{J}_n^m . That is, we have $[\mathfrak{a}] \cdot [\mathfrak{b}] = [\mathfrak{a} \cdot \mathfrak{b}]$ where $[\mathfrak{a} \cdot \mathfrak{b}]$ is zero if it contains less than m vertical arcs. Similarly, we have $[\mathfrak{a}] \circ [\mathfrak{b}] = [\mathfrak{a} \circ \mathfrak{b}]$ in case of “ \circ ”. By abuse of notation we shall identify $[\mathfrak{a}]$ with \mathfrak{a} . Note that \mathbb{I}_n^n is isomorphic to the group algebra $K[S_n]$. Similarly, \mathbb{L}_n has the filtration

$$(1.4) \quad \mathbb{L}_n^0 \subsetneq \mathbb{L}_n^1 \subsetneq \cdots \subsetneq \mathbb{L}_n^{n-1} \subsetneq \mathbb{L}_n^n = \mathbb{L}_n$$

and by abuse of notation we denote the quotients $\mathbb{L}_n^m / \mathbb{L}_n^{m-1}$ and the set all \mathcal{L}_n -diagrams with exactly m vertical arcs again by \mathbb{I}_n^m and \mathcal{J}_n^m , respectively.

An integer partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_s)$, where $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_s$ is a weakly decreasing sequence of positive integers. If $\sum_i \lambda_i = n$, we write $\lambda \vdash n$. Since any irreducible S_n -module is indexed by a partition [12] λ we write them as S^λ . The dimension of S^λ is denoted by f^λ and its character by χ^λ . The integers λ_i is called the parts of λ . The Ferrer diagram associated with a partition λ is a collection of boxes, $[\lambda]$, in \mathbb{Z}^2 using matrix-style coordinates. The boxes are arranged in left-justified rows with weakly decreasing numbers of boxes in each row. For a box $p = (i, j)$ in $[\lambda]$, $j - i$ is the content of p , denoted by $c(p)$. If λ and μ are two partitions such that $\lambda_i \geq \mu_i$ for all i , then we say λ contains μ and write $\mu \subseteq \lambda$. If $\mu \subseteq \lambda$, then the skew partition λ/μ is the set $[\lambda]/[\mu]$. A special case is when λ/μ contains one box only, denoted by $\lambda \sqsupset \mu$. If we identify λ with a Ferrer diagram, then an inner corner of λ is a node $(i, j) \in \lambda$ whose removal leaves the Ferrers diagram of a partition. Any partition μ_1 obtained by such a removal is denoted by $\mu_1 \sqsubset \lambda$. An outer corner of λ is a node $(i, j) \notin \lambda$ whose addition produces the Ferrer diagram of a partition. Any partition μ_2 obtained by such an addition is denoted by $\lambda \sqsubset \mu_2$. Let $\text{res}_{S_{n-1}}^{S_n} S^\lambda$ and $\text{ind}_{S_n}^{S_{n+1}} S^\lambda$ denote the

restriction and the induced representation of S^λ . Then we have [12]

$$(1.5) \quad \text{res}_{S_{n-1}}^{S_n} S^\lambda \cong \bigoplus_{\mu_1 \sqsubset \lambda} S^{\mu_1} \quad \text{and} \quad \text{ind}_{S_n}^{S_{n+1}} S^\lambda \cong \bigoplus_{\lambda \sqsubset \mu_2} S^{\mu_2}.$$

We proceed by describing the induced representation [14] in a specific way. For any $1 \leq j \leq n-t$, set $\tau_j = (j, n+1-t)$ and $\tau_{n+1-t} = 1$. Then $\{\tau_r \mid 1 \leq j \leq n+1-t\}$ is a set of representatives of S_{n+1-t}/S_{n-t} and

$$(1.6) \quad \text{ind}_{S_{n-t}}^{S_{n+1-t}} S^\lambda \cong K[S_{n+1-t}] \otimes_{K[S_{n-t}]} S^\lambda \cong \bigoplus_{j=1}^{n+1-t} (S^\lambda, j).$$

Here, the S_{n+1-t} -action on $\text{ind}_{S_{n-t}}^{S_{n+1-t}} S^\lambda$ is given as follows: for given $\sigma \in S_{n+1-t}$ and $1 \leq j \leq n+1-t$, let s be the unique index such that $\sigma\tau_j \in \tau_s S_{n-t}$ holds, then

$$(1.7) \quad \sigma \cdot (w, j) = ((\tau_s^{-1} \sigma \tau_j)w, s).$$

In the following, let \mathbb{X}_n denote either \mathbb{A}_n or \mathbb{L}_n . Let M be a \mathbb{X}_n -left module. Then $\text{res}_{n-1}(M)$ denotes the \mathbb{X}_{n-1} -left module, obtained via restriction with respect to the embedding $\epsilon_n: \mathbb{X}_{n-1} \longrightarrow \mathbb{X}_n$ and $\text{ind}_{n+1}(M) = \mathbb{X}_{n+1} \otimes_{\mathbb{X}_n} M$ denotes the induced \mathbb{X}_{n+1} -left module.

2. \mathbb{X}_n -MODULES

The semisimplicity of \mathbb{X}_n is closely tied to the structure of \mathbb{X}_n -modules. Therefore we shall begin by establishing their basic properties. The latter are a result of the general machinery derived from the fact that \mathbb{A}_n and \mathbb{L}_n are for $x \neq 0$ quasi-hereditary algebras. However, we shall prove them directly. Let $\mathbf{u}_{n,t}$ denote the diagram having straight verticals except of loops incident to $(n-t+1), \dots, n$ and $(n-t+1)', \dots, n'$, respectively. Pictorially,

$$\mathbf{u}_{n,t} = \begin{array}{ccccccc} & 1 & & n-t & & n-t+1 & n \\ & \bullet & & \bullet & & \bullet & \bullet \\ & \vdots & & \vdots & & \vdots & \vdots \\ & \bullet & & \bullet & & \bullet & \bullet \end{array}$$

Let $x \neq 0$ and $\lambda \vdash (n-t) \leq n$ be a partition, we set

$$(2.1) \quad \mathcal{M}_{\mathbb{X}_n}(\lambda) = \mathbb{I}_n^{n-t} \mathbf{u}_{n,t} \otimes_{S_{n-t}} S^\lambda \quad \text{and} \quad \mathcal{N}_{\mathbb{X}_n}(\lambda) = \{w \in \mathcal{M}_{\mathbb{X}_n}(\lambda) \mid \mathbb{I}_n^{n-t} w = 0\}.$$

$\mathcal{M}_{\mathbb{X}_n}(\lambda)$ and $\mathcal{N}_{\mathbb{X}_n}(\lambda)$ become via linear extension of the action

$$(2.2) \quad \mathbf{b} \cdot (\mathbf{a} \otimes v) = (\mathbf{ba}) \otimes v,$$

\mathbb{X}_n - and \mathbb{I}_n^{n-t} -left modules, respectively. Indeed, for any $0 \leq t \leq n$, $\mathbb{X}_n^{n-t} \triangleleft \mathbb{X}_n$ is a two sided ideal, which implies that $\mathcal{N}_{\mathbb{X}_n}(\lambda)$ is a \mathbb{X}_n -invariant subspace.

Proposition 1. *Let $x \neq 0$ and $\lambda \vdash (n-t) \leq n$ be a partition, then the following assertions hold*
 (a) $\mathcal{M}_{\mathbb{X}_n}(\lambda)/\mathcal{N}_{\mathbb{X}_n}(\lambda)$ *is irreducible as a \mathbb{X}_n -module and \mathbb{I}_n^{n-t} -module, respectively. In particular, $\mathcal{M}_{\mathbb{X}_n}(\lambda)$ is irreducible if and only if $\mathcal{N}_{\mathbb{X}_n}(\lambda) = 0$.*
 (b) $\mathcal{N}_{\mathbb{X}_n}(\lambda)$ *is a maximal \mathbb{X}_n -submodule of $\mathcal{M}_{\mathbb{X}_n}(\lambda)$ and $\mathcal{N}_{\mathbb{X}_n}(\lambda)$ is unique.*
 (c) *For any irreducible \mathbb{X}_n -module, V , there exists a partition $\lambda \vdash m \leq n$ with the property $V \cong \mathcal{M}_{\mathbb{X}_n}(\lambda)/\mathcal{N}_{\mathbb{X}_n}(\lambda)$.*

Proof. We first prove (a). Since $\mathcal{N}_{\mathbb{X}_n}(\lambda)$ is a \mathbb{X}_n -invariant subspace, $\mathcal{M}_{\mathbb{X}_n}(\lambda)/\mathcal{N}_{\mathbb{X}_n}(\lambda)$ is a \mathbb{X}_n - and \mathbb{I}_n^{n-t} -module.

Claim. Any $v \in \mathcal{M}_{\mathbb{X}_n}(\lambda) \setminus \mathcal{N}_{\mathbb{X}_n}(\lambda)$ has the property $\mathbb{I}_n^{n-t}v = \mathcal{M}_{\mathbb{X}_n}(\lambda)$.

To prove the claim we represent $v = \sum_i \mathbf{a}_i \otimes v_i$, where $\mathbf{a}_i \in \mathcal{J}_n^{n-t}$ and $\text{bot}(\mathbf{a}_i) = \text{bot}(\mathbf{u}_{n,t})$. Let $\delta_{\mathbf{b}\mathbf{a}_i} = 1$ if $\mathbf{b}\mathbf{a}_i \neq 0$ and $\delta_{\mathbf{b}\mathbf{a}_i} = 0$ in \mathbb{I}_n^{n-t} , otherwise. For an arbitrary diagram, $\mathbf{b} \in \mathcal{J}_n^{n-t}$, we have

$$(2.3) \quad \mathbf{b} \cdot \sum_i \mathbf{a}_i \otimes v_i = \sum_i (\mathbf{b}\mathbf{a}_i) \otimes v_i = \tilde{\mathbf{b}} \otimes \sum_i \delta_{\mathbf{b}\mathbf{a}_i} x^{\ell(\mathbf{a}_i, \mathbf{b})} \sigma_{\mathbf{a}_i, \mathbf{b}} v_i,$$

where $\ell(\mathbf{b}, \mathbf{a}_i)$ denotes the number of inner components in $G(\mathbf{b}, \mathbf{a}_i)$, $\sigma_{\mathbf{b}, \mathbf{a}_i} \in S_{n-t}$ is such that the diagram $\tilde{\mathbf{b}} \in \mathcal{J}_n^{n-t}$ has noncrossing verticals, has $\text{top}(\tilde{\mathbf{b}}) = \text{top}(\mathbf{b})$ and satisfies

$$(2.4) \quad x^{\ell(\mathbf{b}, \mathbf{a}_i)} \tilde{\mathbf{b}} \sigma_{\mathbf{b}, \mathbf{a}_i} = \mathbf{b}\mathbf{a}_i.$$

For any $v \in \mathcal{M}_{\mathbb{X}_n}(\lambda) \setminus \mathcal{N}_{\mathbb{X}_n}(\lambda)$ we have $\mathbb{I}_n^{n-t}v \neq 0$, whence there exists some $\mathbf{b}_0 \in \mathcal{J}_n^{n-t}$ such that

$$(2.5) \quad \mathbf{b}_0 \cdot \sum_i \mathbf{a}_i \otimes v_i = \sum_i (\mathbf{b}_0 \mathbf{a}_i) \otimes v_i = \tilde{\mathbf{b}}_0 \otimes \sum_i \delta_{\mathbf{b}_0 \mathbf{a}_i} x^{\ell(\mathbf{b}_0, \mathbf{a}_i)} \sigma_{\mathbf{b}_0, \mathbf{a}_i} v_i \neq 0,$$

where $\text{top}(\tilde{\mathbf{b}}_0) = \text{top}(\mathbf{b}_0)$, $\tilde{\mathbf{b}}_0 \in \mathcal{J}_n^{n-t}$ has noncrossing verticals and $\sigma_{\mathbf{b}_0, \mathbf{a}_i} \in S_{n-t}$ is such that

$$(2.6) \quad x^{\ell(\mathbf{b}_0, \mathbf{a}_i)} \tilde{\mathbf{b}}_0 \sigma_{\mathbf{b}_0, \mathbf{a}_i} = \mathbf{b}_0 \mathbf{a}_i.$$

For arbitrary $\mathbf{b} \in \mathcal{J}_n^{n-t}$ we consider the element \mathbf{b}^\dagger having the properties: $\text{top}(\mathbf{b}^\dagger) = \text{top}(\mathbf{b})$, $\text{bot}(\mathbf{b}^\dagger) = \text{bot}(\mathbf{b}_0)$, having $n-t$ vertical arcs and satisfying

$$(2.7) \quad \mathbf{b}^\dagger \mathbf{a}_i = x^{\ell(\mathbf{b}_0, \mathbf{a}_i)} \tilde{\mathbf{b}}^\dagger \sigma_{\mathbf{b}_0, \mathbf{a}_i},$$

where $\tilde{\mathbf{b}}^\dagger \in \mathcal{J}_n^{n-t}$ has noncrossing verticals and $\text{top}(\tilde{\mathbf{b}}^\dagger) = \text{top}(\mathbf{b}^\dagger) = \text{top}(\mathbf{b})$. Multiplying with \mathbf{b}^\dagger we obtain

$$\mathbf{b}^\dagger \cdot \sum_i \mathbf{a}_i \otimes v_i = \sum_i (\mathbf{b}^\dagger \mathbf{a}_i) \otimes v_i = \tilde{\mathbf{b}}^\dagger \otimes \sum_i \delta_{\mathbf{b}_0 \mathbf{a}_i} x^{\ell(\mathbf{b}_0, \mathbf{a}_i)} \sigma_{\mathbf{b}_0, \mathbf{a}_i} v_i \neq 0.$$

We set $w = \sum_i \delta_{\mathbf{b}_0 \mathbf{a}_i} x^{\ell(\mathbf{b}_0, \mathbf{a}_i)} \sigma_{\mathbf{b}_0, \mathbf{a}_i} v_i$ and note that $w \neq 0$ holds. Since S^λ is irreducible, for any $0 \neq u$ the elements $\sigma_0 u$, $\sigma_0 \in S_{n-t}$ generate S^λ . Since for any $\sigma_0 \in S_{n-t}$ there exists some $g(\sigma_0, \mathbf{b}^\dagger) \in \mathbb{I}_n^{n-t}$ with the property

$$(2.8) \quad g(\sigma_0, \mathbf{b}^\dagger) \cdot \tilde{\mathbf{b}}^\dagger = x^m \tilde{\mathbf{b}}^\dagger \sigma_0 \quad \text{for some } m \in \mathbb{Z},$$

we conclude

$$(2.9) \quad g(\sigma_0) \cdot \mathbf{b}^\dagger \cdot \sum_i \mathbf{a}_i \otimes v_i = g(\sigma_0) \cdot \tilde{\mathbf{b}}^\dagger \otimes w = x^m \tilde{\mathbf{b}}^\dagger \otimes \sigma_0 w.$$

Accordingly, $\mathbb{I}_n^{n-t} \cdot v = \mathbb{I}_n^{n-t} \mathbf{u}_{n,t} \otimes_{S_{n-t}} S^\lambda$ and the Claim is proved.

As a result, any nontrivial $\mathcal{M}_{\mathbb{X}_n}(\lambda)/\mathcal{N}_{\mathbb{X}_n}(\lambda)$ -element generates $\mathcal{M}_{\mathbb{X}_n}(\lambda)/\mathcal{N}_{\mathbb{X}_n}(\lambda)$, which is equivalent to $\mathcal{M}_{\mathbb{X}_n}(\lambda)/\mathcal{N}_{\mathbb{X}_n}(\lambda)$ being an irreducible \mathbb{I}_n^{n-t} -left module. This action extends to a unique \mathbb{X}_n -left action with respect to which $\mathcal{M}_{\mathbb{X}_n}(\lambda)/\mathcal{N}_{\mathbb{X}_n}(\lambda)$ is an irreducible \mathbb{X}_n -module. This proves assertion (a).

We next prove (b): the maximality of $\mathcal{N}_{\mathbb{X}_n}(\lambda)$ follows from the irreducibility of $\mathcal{M}_{\mathbb{X}_n}(\lambda)/\mathcal{N}_{\mathbb{X}_n}(\lambda)$. It remains to show that $\mathcal{N}_{\mathbb{X}_n}(\lambda)$ is unique. For this purpose, let M be a maximal \mathbb{X}_n -left submodule of $\mathcal{M}_{\mathbb{X}_n}(\lambda)$ different from $\mathcal{N}_{\mathbb{X}_n}(\lambda)$. Then there exist a $v \in M \setminus \mathcal{N}_{\mathbb{X}_n}(\lambda)$, which, according to (a) generates $\mathcal{M}_{\mathbb{X}_n}(\lambda)$. Consequently, any maximal $\mathcal{M}_{\mathbb{X}_n}(\lambda)$ -module, different from $\mathcal{N}_{\mathbb{X}_n}(\lambda)$, is equal to $\mathcal{M}_{\mathbb{X}_n}(\lambda)$ and (b) follows.

Next we show (c). Let $(n-t)$ be the smallest integer with the property \mathbb{X}_n^{n-t} is not acting trivially on V . Consider the set $V_0 = \{v \in V \mid \mathbb{I}_n^{n-t} v = 0\}$. Clearly, since $\mathbb{X}_n^{n-t} \triangleleft \mathbb{X}_n$ is a two sided ideal, V_0 is an \mathbb{X}_n -invariant subspace and the irreducibility of V implies either $V_0 = 0$ or $V_0 = V$. By definition of $(n-t)$, there exists a $v \in V$ such that $\mathbb{I}_n^{n-t} v \neq 0$, whence $V_0 = 0$. Therefore, any $0 \neq v \in V$ has the property $\mathbb{I}_n^{n-t} v \neq 0$ and $\mathbb{I}_n^{n-t} v$ is \mathbb{A}_n -invariant. Since V is an irreducible \mathbb{X}_n -module we have $\mathbb{I}_n^{n-t} v = V$. Accordingly, V is also an irreducible \mathbb{I}_n^{n-t} -left module.

As an \mathbb{I}_n^{n-t} -left module the algebra \mathbb{I}_n^{n-t} decomposes into a direct sum of modules that are isomorphic to $\mathcal{M}_{\mathbb{X}_n}(\lambda)$, for $\lambda \vdash (n-t)$, i.e.

$$(2.10) \quad \mathbb{I}_n^{n-t} \cong \bigoplus_{\lambda \vdash (n-t)} n_\lambda \mathcal{M}_{\mathbb{X}_n}(\lambda),$$

where n_λ denotes the multiplicity of $\mathcal{M}_{\mathbb{X}_n}(\lambda)$ in \mathbb{I}_n^{n-t} . Clearly we have for any $0 \neq v \in V$ the surjective morphism of \mathbb{I}_n^{n-t} -left modules $\phi_v: \mathbb{I}_n^{n-t} \longrightarrow V$, given by $\mathbf{a} \mapsto \mathbf{a} \cdot v$. Accordingly there exists a partition $\lambda \vdash (n-t)$ and a surjective morphism of \mathbb{I}_n^{n-t} -left modules induced by ϕ_v :

$$\phi_v^\lambda: \mathcal{M}_{\mathbb{X}_n}(\lambda) \longrightarrow V.$$

Assertion (a) and (b) imply $\ker(\phi_v^\lambda) = \mathcal{N}_{\mathbb{X}_n}(\lambda)$, i.e. we have $\mathcal{M}_{\mathbb{X}_n}(\lambda)/\mathcal{N}_{\mathbb{X}_n}(\lambda) \cong V$ and the proof of Proposition 1 is complete. \square

The next result connects semisimplicity of \mathbb{X}_n with the existence of nontrivial morphisms between the modules $\mathcal{M}_{\mathbb{X}_n}(\lambda)$ and $\mathcal{M}_{\mathbb{X}_n}(\mu)$. Indeed, if \mathbb{X}_n is not semisimple, then there exists some module $\mathcal{M}_{\mathbb{X}_n}(\mu)$, $\mu \vdash m < n$ with a nontrivial maximal submodule $\mathcal{N}_{\mathbb{X}_n}(\mu)$. In the following we denote by $\text{Rad}(\mathbb{X}_n)$ the Jacobson radical of \mathbb{X}_n , i.e. \mathbb{X}_n is semisimple if and only if $\text{Rad}(\mathbb{X}_n) = 0$.

Proposition 2. *If \mathbb{X}_n is not semisimple, then there exist two partitions μ, λ , where $|\mu| < |\lambda| \leq n$ and a short exact sequence of \mathbb{X}_n -modules*

$$(2.11) \quad 0 \longrightarrow \mathcal{N}_{\mathbb{X}_n}(\lambda) \longrightarrow \mathcal{M}_{\mathbb{X}_n}(\lambda) \xrightarrow{\varphi_n} \mathcal{M}_{\mathbb{X}_n}(\mu).$$

Proof. Suppose first that $\mathcal{M}_{\mathbb{X}_n}(\mu)$ is for any partition $\mu \vdash m$, $m < n$, irreducible. We claim that \mathbb{X}_n is in this case semisimple. To this end we observe that for $\mu \vdash n$, we have $\mathcal{M}_{\mathbb{X}_n}(\mu) \cong S^\mu$, i.e. for arbitrary partition μ , the module $\mathcal{M}_{\mathbb{X}_n}(\mu)$ is irreducible. In view of

$$\mathbb{I}_n^m \cong \bigoplus_{\mu \vdash m} n_\mu \mathcal{M}_{\mathbb{X}_n}(\mu),$$

for any $0 \leq m \leq n$, the F -algebras $\mathbb{X}_n^m / \mathbb{X}_n^{m-1} \cong \mathbb{I}_n^m$ and in particular $\mathbb{X}_n^0 \cong \mathbb{I}_n^0$, are semisimple. Since $\text{Rad}(\mathbb{X}_n^m)$ is a nilpotent ideal so is $(\text{Rad}(\mathbb{X}_n^m) + \mathbb{X}_n^{m-1}) / \mathbb{X}_n^{m-1}$ and we obtain

$$(\text{Rad}(\mathbb{X}_n^m) + \mathbb{X}_n^{m-1}) / \mathbb{X}_n^{m-1} \subset \text{Rad}(\mathbb{X}_n^m / \mathbb{X}_n^{m-1}) = 0.$$

We next observe using $\text{Rad}(\mathbb{X}_n^m) \cap \mathbb{X}_n^{m-1} = \text{Rad}(\mathbb{X}_n^{m-1})$

$$\begin{aligned} (\text{Rad}(\mathbb{X}_n^m) + \mathbb{X}_n^{m-1}) / \mathbb{X}_n^{m-1} &\cong \text{Rad}(\mathbb{X}_n^m) / (\text{Rad}(\mathbb{X}_n^m) \cap \mathbb{X}_n^{m-1}) \\ &\cong \text{Rad}(\mathbb{X}_n^m) / \text{Rad}(\mathbb{X}_n^{m-1}). \end{aligned}$$

Consequently we have for $1 \leq m \leq n$ the inclusion $\text{Rad}(\mathbb{X}_n^m) \subset \text{Rad}(\mathbb{X}_n^{m-1})$, which implies $\text{Rad}(\mathbb{X}_n) \subset \text{Rad}(\mathbb{X}_n^0) = 0$, i.e. \mathbb{X}_n is semisimple.

Thus, if \mathbb{X}_n is not semisimple, there exists a partition $\mu \vdash m$, $m < n$, such that $\mathcal{M}_{\mathbb{X}_n}(\mu)$ is not irreducible. Then there exists according to Proposition 1, assertion (b), the nontrivial, maximal submodule $\mathcal{N}_{\mathbb{X}_n}(\mu) \subset \mathcal{M}_{\mathbb{X}_n}(\mu)$. Let m_0 be the smallest integer such that $\mathbb{X}_n^{m_0}$ acts nontrivially on $\mathcal{N}_{\mathbb{X}_n}(\mu)$. By definition we have for any $v \in \mathcal{N}_{\mathbb{X}_n}(\mu)$, $\mathbb{I}_n^m v = 0$, whence $m < m_0$. $\mathcal{N}_{\mathbb{X}_n}(\mu)$ is then a nontrivial $\mathbb{I}_n^{m_0}$ -left module and there exists an irreducible $\mathbb{I}_n^{m_0}$ -submodule $W \subset \mathcal{N}_{\mathbb{X}_n}(\mu)$. According to Proposition 1, assertion (c), W is isomorphic to $\mathcal{M}_{\mathbb{X}_n}(\lambda) / \mathcal{N}_{\mathbb{X}_n}(\lambda)$ for some $\lambda \vdash m_0$, i.e. $|\mu| < |\lambda| \leq n$. Therefore there exists a partition λ and a nontrivial morphism of \mathbb{X}_n -modules $\varphi_n: \mathcal{M}_{\mathbb{X}_n}(\lambda) \longrightarrow \mathcal{M}_{\mathbb{X}_n}(\mu)$, such that $\ker(\varphi_n) = \mathcal{N}_{\mathbb{X}_n}(\lambda)$ and $|\mu| < |\lambda| \leq n$ and the proposition follows. \square

3. RESTRICTION AND INDUCTION

We shall begin by showing that \mathbb{A}_n has the generators $\mathfrak{g}_{i-1}, \mathfrak{e}_{i-1}, \mathfrak{u}_j$, $2 \leq i \leq n$, $1 \leq j \leq n$.

Lemma 1. *Any diagram $\mathfrak{a} \in \mathcal{A}_n$ is either contained in \mathcal{A}_{n-1} or of the form*


$$(3.1) \quad \mathfrak{a} = \mathfrak{a}' \mathfrak{x} \mathfrak{b}', \quad \mathfrak{a}', \mathfrak{b}' \in \mathcal{A}_{n-1}, \quad \mathfrak{x} \in \{\mathfrak{g}_{n-1}, \mathfrak{e}_{n-1}, \mathfrak{u}_n\}.$$

In particular, we have $\mathbb{A}_n = \langle S_n, \mathfrak{e}_{n-1}, \mathfrak{u}_n \rangle$ and $\mathbb{L}_n = \langle S_n, \mathfrak{u}_n \rangle$.

Proof. Any diagram not contained in \mathcal{A}_{n-1} has either (a) none or two loops at the vertices n, n' , (b) exactly one loop over $n(n')$ and at least one loop over some vertex $i'(i)$, where $i < n$ or (c) exactly one loop over $n(n')$ and no loops over $i'(i)$, where $i < n$. From this we derive

$$(3.2) \quad \mathfrak{a} = \mathfrak{a}' \mathfrak{y} \mathfrak{b}' \quad \text{where} \quad \mathfrak{a}', \mathfrak{b}' \in \mathcal{A}_{n-1} \quad \text{and} \quad \mathfrak{y} \in \begin{cases} \{\mathfrak{g}_{n-1}, \mathfrak{e}_{n-1}, \mathfrak{u}_n\} & \text{(a)} \\ \{\mathfrak{d}_1, \mathfrak{d}_1^*, \mathfrak{d}_2, \mathfrak{d}_2^*\} & \text{(b)} \\ \{\mathfrak{d}_3, \mathfrak{d}_3^*, \mathfrak{d}_4, \mathfrak{d}_4^*\} & \text{(c)} \end{cases}$$

where




We can express the diagrams $\mathfrak{d}_1, \dots, \mathfrak{d}_4$ via the generators \mathfrak{g}_i , \mathfrak{e}_i and \mathfrak{u}_i as follows

$$\mathfrak{d}_1 = \mathfrak{e}_{n-2} \mathfrak{u}_{n-1} \mathfrak{u}_{n-2} \mathfrak{e}_{n-1}, \quad \mathfrak{d}_2 = \mathfrak{u}_n \mathfrak{g}_{n-1}, \quad \mathfrak{d}_3 = \mathfrak{u}_n \mathfrak{e}_{n-1} \mathfrak{e}_{n-2}, \quad \mathfrak{d}_4 = \mathfrak{u}_{n-1} \mathfrak{e}_{n-1}.$$

We next observe that the relations

$$\mathfrak{u}_n \mathfrak{e}_{n-1} = \mathfrak{u}_{n-1} \mathfrak{e}_{n-1} \quad \mathfrak{g}_{n-1} \mathfrak{u}_{n-1} = \mathfrak{u}_n \mathfrak{g}_{n-1} \quad \mathfrak{u}_{n-1} \mathfrak{g}_{n-1} = \mathfrak{g}_{n-1} \mathfrak{u}_n$$



imply eq. (3.2) from which the lemma follows. \square

The next theorem analyzes the restriction in \mathbb{A}_n and follows the ideas of Doran *et al.* [4] in the case of \mathbb{B}_n . We find the following new phenomenon for \mathbb{A}_n : for $\lambda \vdash (n-t)$, where $t \geq 1$, there exists an embedding of $\mathcal{M}_{\mathbb{A}_{n-1}}(\lambda)$ into $\text{res}_{n-1}(\mathcal{M}_{\mathbb{A}_n}(\lambda))$. Such an embedding does not exist for \mathbb{B}_n . We shall employ it in Lemma 2 in order to show that if $\text{hom}_{\mathbb{X}_n}(\mathcal{M}_{\mathbb{X}_n}(\lambda), \mathcal{M}_{\mathbb{X}_n}(\mu)) \neq 0$ then we can, without loss of generality, assume that $\lambda \vdash n$.

Theorem 1. *Let $n, t \in \mathbb{N}$ and $\lambda \vdash (n-t)$ where $1 \leq t \leq n$. Then there exists the exact sequence of \mathbb{A}_{n-1} -modules*


$$(3.3) \quad 0 \longrightarrow \bigoplus_{\alpha \sqsubseteq \lambda} \mathcal{M}_{\mathbb{A}_{n-1}}(\alpha) \longrightarrow \text{res}_{n-1}(\mathcal{M}_{\mathbb{A}_n}(\lambda)) \longrightarrow \bigoplus_{\lambda \sqsubset \beta} \mathcal{M}_{\mathbb{A}_{n-1}}(\beta) \longrightarrow 0.$$

Proof. Claim 1. There exists the following short exact sequence of \mathbb{A}_{n-1} -left modules

$$(3.4) \quad 0 \longrightarrow \bigoplus_{\alpha \sqsubseteq \lambda} \mathcal{M}_{\mathbb{A}_{n-1}}(\alpha) \longrightarrow \text{res}_{n-1}(\mathcal{M}_{\mathbb{A}_n}(\lambda)).$$

Let $F_n^1(\lambda)$ denote the $\mathcal{M}_{\mathbb{A}_n}(\lambda)$ -subspace generated by all tensors $\mathbf{a} \otimes w$, where \mathbf{a} is a $\mathbb{I}_n^{n-t} \mathbf{u}_{n,t}$ -diagram in which all vertical edges are noncrossing and the top-vertex n is incident to a vertical edge. Obviously, any tensor $\mathbf{b} \otimes w \in \mathbb{I}_n^{n-t} \mathbf{u}_{n,t} \otimes_{S_{n-t}} S^\lambda$ in which n is incident to a vertical edge, satisfies $\mathbf{b} \otimes w = \mathbf{a} \otimes \sigma w$ for some $\sigma \in S_{n-t}$. Let $f_1(\mathbf{a})$ be the diagram derived from \mathbf{a} by removing n and $(n-t)'$ and by shifting all bottom vertices $\ell' > (n-t)'$ down by one. f_1 induces the mapping

$$(3.5) \quad \begin{aligned} \varphi_1: F_n^1(\lambda) &\longrightarrow \mathbb{I}_{n-1}^{n-1-t} \mathbf{u}_{n-1,t} \otimes_{S_{(n-1)-t}} \text{res}_{S_{n-1-t}}(S^\lambda) \\ \mathbf{a} \otimes w &\longmapsto f_1(\mathbf{a}) \otimes w. \end{aligned}$$



We next prove that φ_1 is bijective. Indeed, for any $\mathbb{I}_{n-1}^{n-1-t} \mathbf{u}_{n-1,t}$ -diagram, \mathfrak{x} , there exists a unique permutation $\sigma_0 \in S_{n-1-t}$ such that the vertical edges in $\mathfrak{x}\sigma_0$ are noncrossing. Furthermore we have $\mathfrak{x} \otimes w = \mathfrak{x}\sigma_0 \otimes \sigma_0^{-1}w$. Clearly, the tensor $\mathfrak{x}\sigma_0 \otimes \sigma_0^{-1}w$ has a unique φ_1 -preimage, $f_1^{-1}(\mathfrak{x}\sigma_0) \otimes \sigma_0^{-1}w$ where $f_1^{-1}(\mathfrak{x}\sigma_0)$ is obtained by shifting the bottom vertices $\ell' \geq (n-t)'$ up by one and by adding the vertices n and $(n-t)'$ together with an vertical edge connecting them. This proves that φ_1 is bijective.

We next show that $F_n^1(\lambda)$ is, via the natural embedding $\epsilon_n: \mathbb{A}_{n-1} \longrightarrow \mathbb{A}_n$, an \mathbb{A}_{n-1} -module. In view of Lemma 1 it suffices to show

$$\mathfrak{x} \cdot (\mathbf{a} \otimes v_i) \in F_n^1(\lambda),$$

where $\mathfrak{x} \in \{\sigma, \mathbf{e}_i, \mathbf{u}_j\}$, $1 \leq j \leq n-1$, $1 \leq i \leq n-2$ and $\sigma \in S_{n-1}$. Let \mathbf{a} be a $\mathbb{I}_n^{n-t} \mathbf{u}_{n,t}$ -diagram in which all vertical edges are noncrossing and the top-vertex n is incident to a vertical edge and let $\sigma \in S_{n-1}$. Then there exist a unique $\mathbb{I}_n^{n-t} \mathbf{u}_{n,t}$ -diagram, \mathbf{a}' , with noncrossing vertical edges, in which n is connected to $(n-t)'$ and a permutation $\sigma_0 \in S_{(n-1)-t}$ such that $\sigma\mathbf{a} = \mathbf{a}'\sigma_0$ holds. Consequently,

$$\sigma \cdot (\mathbf{a} \otimes w) = \mathbf{a}'\sigma_0 \otimes w = \mathbf{a}' \otimes \sigma_0 w,$$

i.e. $\sigma \cdot (\mathbf{a} \otimes w) \in F_n^1(\lambda)$. The cases $\mathbf{e}_i \cdot \mathbf{a} \otimes v_j$ and $\mathbf{u}_{i+1} \cdot \mathbf{a} \otimes v_j$ follow analogously. We next show that φ_1 is an isomorphism of \mathbb{A}_{n-1} -modules, that is we prove $\mathbf{b} \cdot \varphi_1(\zeta) = \varphi_1(\mathbf{b} \cdot \zeta)$. Indeed, for $\mathfrak{x} \in \{\sigma, \mathbf{e}_i, \mathbf{u}_j\}$


$$\mathfrak{x} \cdot (f(\mathbf{a}) \otimes w) = f(\mathfrak{x}\mathbf{a}) \otimes w,$$

since neither vertex n or its incident bottom vertex $(n-t)'$ are affected by left multiplication with the elements $\sigma, \mathbf{e}_i, \mathbf{u}_j$.

Let $F_n^2(\lambda) \subset \mathcal{M}_{\mathbb{A}_n}(\lambda)$ be the subspace generated by all tensors $\mathbf{a} \otimes v_i$, where $\mathbf{a} \in \mathbb{I}_n^{n-t} \mathbf{u}_{n,t}$ is a diagram having a loop at vertex n . Let $f_2(\mathbf{a}) \in \mathbb{I}_{n-1}^{n-t} \mathbf{u}_{n-1,t-1}$ be the diagram obtained by removing the vertices n and n' together with their loops. It is straightforward to show that f_2 induces the isomorphism of \mathbb{A}_{n-1} -modules

$$(3.6) \quad \begin{aligned} \varphi_2: F_n^2(\lambda) &\longrightarrow \mathbb{I}_{n-1}^{n-t} \mathbf{u}_{n-1,t-1} \otimes_{S_{n-t}} S^\lambda \\ \mathbf{a} \otimes w &\longmapsto f_2(\mathbf{a}) \otimes w, \end{aligned}$$

where $\mathbb{I}_{n-1}^{n-t} \mathbf{u}_{n-1,t-1} \otimes_{S_{n-t}} S^\lambda \cong \mathcal{M}_{\mathbb{A}_{n-1}}(\lambda)$.



In view of $\text{res}_{S_{n-1-t}}(S^\lambda) \cong \bigoplus_{\alpha \sqsubset \lambda} S^\alpha$ we derive

$$\begin{aligned} F_n^1(\lambda) \oplus F_n^2(\lambda) &\cong [\mathbb{I}_{n-1}^{n-1-t} \mathbf{u}_{n-1,t} \otimes_{S_{(n-1)-t}} \text{res}_{S_{n-1-t}}(S^\lambda)] \oplus [\mathbb{I}_{n-1}^{n-t} \mathbf{u}_{n-1,t-1} \otimes_{S_{n-t}} S^\lambda] \\ &\cong \bigoplus_{\alpha \sqsubset \lambda} [\mathbb{I}_{n-1}^{n-1-t} \mathbf{u}_{n-1,t} \otimes_{S_{(n-1)-t}} S^\alpha] \oplus [\mathbb{I}_{n-1}^{n-t} \mathbf{u}_{n-1,t-1} \otimes_{S_{n-t}} S^\lambda], \end{aligned}$$


which gives rise to the short exact sequence $0 \longrightarrow \bigoplus_{\alpha \sqsubseteq \lambda} \mathcal{M}_{\mathbb{A}_{n-1}}(\alpha) \longrightarrow \text{res}_{n-1}(\mathcal{M}_{\mathbb{A}_n}(\lambda))$ and Claim 1 follows.

Claim 2. Let $F_n(\lambda) = F_n^1(\lambda) \oplus F_n^2(\lambda)$, then we have an isomorphism of \mathbb{A}_{n-1} -left modules

$$(3.7) \quad \text{res}_{n-1}(\mathcal{M}_{\mathbb{A}_n}(\lambda)/F_n(\lambda)) \cong \bigoplus_{\lambda \sqsubset \beta} \mathcal{M}_{\mathbb{A}_{n-1}}(\beta).$$

Let $G_n(\lambda)$ denote the space generated by all tensors of the form $\mathbf{c} \otimes w$, where $\mathbf{c} \in \mathbb{I}_n^{n-t} \mathbf{u}_{n,t}$ is a diagram with noncrossing vertical arcs and a horizontal arc incident to n . Let $f_3(\mathbf{c})$ be the diagram obtained from \mathbf{c} as follows: one removes n together with its incident horizontal arc and the bottom-vertex n' together with its incident loop. This leaves a unique top-vertex, r , isolated. Next one removes the loop of the bottom-vertex $(n-t+1)'$ and connects it to r via a vertical arc. We next show that f_3 induces the bijection

$$(3.8) \quad \begin{aligned} \varphi_3: \text{res}_{n-1}(\mathcal{M}_{\mathbb{A}_n}(\lambda)/F_n(\lambda)) &\longrightarrow \mathbb{I}_{n-1}^{n+1-t} \mathbf{u}_{n-1,t-2} \otimes_{S_{n+1-t}} \text{ind}_{S_{n-t}}^{S_{n+1-t}}(S^\lambda) \\ \mathbf{c} \otimes w &\longmapsto f_3(\mathbf{c}) \otimes (w, n+1-t). \end{aligned}$$



Recall that for any $1 \leq j \leq n-t$, $\tau_j = (j, n+1-t)$ and $\tau_{n+1-t} = 1$. Then $S_{n+1-t} = \dot{\bigcup} \tau_j S_{n-t}$, i.e. the τ_j form a set of representatives of S_{n+1-t}/S_{n-t} . We inspect that there exists some $\sigma \in S_{n-t+1}$ such that $f_3(\mathbf{c})\sigma^{-1} = \tilde{\mathbf{c}}$ has noncrossing vertical arcs. Then we have $\sigma = \tau_j \sigma_0$, for some $\sigma_0 \in S_{n-t}$. Therefore, in view of $f_3(\mathbf{c})\sigma^{-1} = \tilde{\mathbf{c}}$, each $f_3(\mathbf{c})$ gives rise to some unique τ_j . Using eq. (1.7) we obtain

$$\begin{aligned} f_3(\mathbf{c})\sigma^{-1} \otimes (w, n+1-t) &= \tilde{\mathbf{c}} \tau_j \sigma_0 \otimes (w, n+1-t) \\ &= \tilde{\mathbf{c}} \tau_j \otimes (\sigma_0 w, n+1-t) \\ &= \tilde{\mathbf{c}} \otimes (\sigma_0 w, j). \end{aligned}$$

There exist exactly $(n+1-t)$ different \mathcal{J}_n^{n-t} -diagrams $\mathbf{c}_1, \dots, \mathbf{c}_{n+1-t}$ having noncrossing vertical arcs in which n is connected to a top-vertex and $\text{bot}(\mathbf{c}_j) = \text{bot}(\mathbf{u}_{n,t})$ with the property

$$(3.9) \quad f_3(\mathbf{c}_j)\sigma^{(j)} = \tilde{\mathbf{c}}$$

for some $\sigma^{(j)} \in S_{n-t+1}$. Since $\dim[\text{ind}_{S_{n-t}}^{S_{n+1-t}}(S^\lambda)] = (n+1-t) \cdot \dim[S^\lambda]$, we obtain

$$(3.10) \quad \dim[\text{res}_{n-1}(\mathcal{M}_{\mathbb{A}_n}(\lambda)/F_n(\lambda))] = \dim\left[\mathbb{I}_{n-1}^{n+1-t} \mathbf{u}_{n-1,t-2} \otimes_{S_{n+1-t}} \text{ind}_{S_{n-t}}^{S_{n+1-t}}(S^\lambda)\right].$$

Therefore it suffices to prove that φ_3 is surjective. $\mathbb{I}_{n-1}^{n+1-t} \mathbf{u}_{n-1,t-2} \otimes_{S_{n+1-t}} \text{ind}_{S_{n-t}}^{S_{n+1-t}}(S^\lambda)$ is generated by tensors of the form $\mathfrak{d} \otimes (w, j)$, where $1 \leq j \leq n+1-t$, $\mathfrak{d} \in \mathcal{J}_{n-1}^{n-t+1}$ with noncrossing vertical arcs, $\text{bot}(\mathfrak{d}) = \text{bot}(\mathbf{u}_{n-1,t-2})$ and $w \in S^\lambda$. Since for $1 \leq j \leq n+1-t$, we have $\tau_j \cdot (w, n+1-t) = (w, j)$ we obtain

$$(3.11) \quad \mathfrak{d} \otimes (w, j) = \mathfrak{d} \otimes \tau_j \cdot (w, n+1-t) = \mathfrak{d}\tau_j \otimes (w, n+1-t).$$

By construction $\mathfrak{d}\tau_j$ is a diagram in which $(n+1-t)'$ connected to a top vertex, which we denote by r . Then there exists some $\sigma_0 \in S_{n-t}$ such that in $\mathfrak{d}\tau_j\sigma_0$ any pair of crossing verticals contains the vertical arc $((n+1-t)', r)$. Let $\mathbf{c} \in \mathcal{J}_n^{n-t}$, be derived from $\mathfrak{d}\tau_j\sigma_0$ by removing $(r, (n+1-t)'),$ adding the vertices n and n' , the loops at $(n+1-t)'$ and n' , as well as the horizontal arc (r, n) . By construction $\text{bot}(\mathbf{c}) = \text{bot}(\mathbf{u}_{n,t})$, \mathbf{c} has noncrossing verticals and we have

$$(3.12) \quad (\mathfrak{d}\tau_j)\sigma_0 = f_3(\mathbf{c}).$$

Consequently, using the fact that the tensor product is over S_{n+1-t}

$$\begin{aligned} \mathfrak{d} \otimes (w, j) &= \mathfrak{d}\tau_j \otimes (w, n+1-t) \\ &= f_3(\mathbf{c})\sigma_0^{-1} \otimes (w, n+1-t) \\ &= f_3(\mathbf{c}) \otimes (\sigma_0^{-1}w, n+1-t), \end{aligned}$$

which proves that φ_3 is surjective. We proceed by showing that φ_3 is an isomorphism of \mathbb{A}_{n-1} -modules. Since any $\sigma \in S_{n-1}$ fixes n we inspect

$$(3.13) \quad \forall \sigma \in S_{n-1}; \quad \varphi_3(\sigma \cdot \mathbf{c} \otimes w) = \sigma \cdot f_3(\mathbf{c}) \otimes (w, n+1-t) = \sigma \cdot \varphi_3(\mathbf{c} \otimes w).$$

We next consider the action of \mathbf{e}_i , $1 \leq i \leq n-2$. Suppose n is connected to r in \mathbf{c} and $r \neq i+1, i$. Then we immediately obtain

$$(3.14) \quad \varphi_3(\mathbf{e}_i \cdot \mathbf{c} \otimes w) = \mathbf{e}_i \cdot f_3(\mathbf{c}) \otimes (w, n+1-t) = \mathbf{e}_i \cdot \varphi_3(\mathbf{c} \otimes w).$$

Without loss of generality we may assume $r = i$. We distinguish three cases:

(1) if $i+1$ is incident to a vertical arc, in $\mathbf{e}_i\mathbf{c}$ the top-vertex n is connected to a bottom vertex, whence $\mathbf{e}_i \cdot \mathbf{c} \otimes w \equiv 0$ modulo $F_n(\lambda)$,

$$f_3 \left(\begin{array}{c} i \quad i+1 \quad n \\ \text{diagram} \end{array} \right) = f_3 \left(\begin{array}{c} i \quad i+1 \quad n \\ \text{diagram} \end{array} \right) \epsilon_i \cdot f_3 \left(\begin{array}{c} i \quad i+1 \quad n \\ \text{diagram} \end{array} \right) = \begin{array}{c} i \quad i+1 \\ \text{diagram} \end{array}$$

On the other hand, in $f_3(\mathbf{c})$, $i+1$ and i are connected to vertical arcs, whence $\epsilon_i \cdot \varphi_3(\mathbf{c} \otimes w)$ has fewer than $(n+1-t)$ vertical arcs and is consequently zero in $\mathbb{I}_{n-1}^{n+1-t} \mathbf{u}_{n-1, t-2}$.

(2) if $i+1$ is incident to a loop, n is incident to a loop in $\epsilon_i \mathbf{c}$. Clearly we then have $f_3(\epsilon_i \mathbf{c}) = \epsilon_i f_3(\mathbf{c})$ implying

$$\varphi_3(\epsilon_i \cdot \mathbf{c} \otimes w) = \epsilon_i \cdot f_3(\mathbf{c}) \otimes (w, n+1-t) = \epsilon_i \cdot \varphi_3(\mathbf{c} \otimes w).$$

$$f_3 \left(\begin{array}{c} i \quad i+1 \quad n \\ \text{diagram} \end{array} \right) = f_3 \left(\begin{array}{c} i \quad i+1 \quad n \\ \text{diagram} \end{array} \right) \epsilon_i \cdot f_3 \left(\begin{array}{c} i \quad i+1 \quad n \\ \text{diagram} \end{array} \right) = \begin{array}{c} i \quad i+1 \\ \text{diagram} \end{array}$$

(3) if $i+1$ is incident to j via a horizontal arc, n is connected to j in $\epsilon_i \mathbf{c}$. Clearly we then have $f_3(\epsilon_i \mathbf{c}) = \epsilon_i f_3(\mathbf{c})$ implying

$$\varphi_3(\epsilon_i \cdot \mathbf{c} \otimes w) = \epsilon_i \cdot f_3(\mathbf{c}) \otimes (w, n+1-t) = \epsilon_i \cdot \varphi_3(\mathbf{c} \otimes w).$$

$$f_3 \left(\begin{array}{c} j \quad i \quad i+1 \quad n \\ \text{diagram} \end{array} \right) = f_3 \left(\begin{array}{c} i \quad i+1 \quad n \\ \text{diagram} \end{array} \right) \epsilon_i \cdot f_3 \left(\begin{array}{c} i \quad i+1 \quad n \\ \text{diagram} \end{array} \right) = \begin{array}{c} j \quad i \quad i+1 \\ \text{diagram} \end{array}$$

Finally we consider the action of u_i , $1 \leq i \leq n-1$. Suppose first $r \neq i$. By definition of f_3 , a vertex $i \neq r$ is in \mathbf{c} incident to a vertical arc if and only if this holds for $f_3(\mathbf{c})$. In this case we have $u_i \mathbf{c} \equiv 0 \pmod{F_n(\lambda)}$ and $u_i f_3(\mathbf{c}) \equiv 0$ in $\mathbb{I}_{n-1}^{n+1-t} \mathbf{u}_{n-1, t-2}$. If i is incident to a loop we have $u_i \mathbf{c} = x\mathbf{c}$ and $u_i f_3(\mathbf{c}) = x f_3(\mathbf{c})$, i.e. $\varphi_3(u_i \cdot \mathbf{c} \otimes w) = u_i \cdot \varphi_3(\mathbf{c} \otimes w)$. Finally, if i is incident to a horizontal arc we have $f_3(u_i \mathbf{c}) = u_i f_3(\mathbf{c})$. Second let $r = i$. On the one hand we obtain $u_i \mathbf{c} \equiv 0$ modulo $F_n(\lambda)$, since the i' -loop of u_i traces back to the top vertex n of $u_i \mathbf{c}$. On the other hand, in $u_i f_3(\mathbf{c})$ the i' -loop of u_i traces back to the bottom vertex $(n+1-t)'$. Consequently, $u_i f_3(\mathbf{c})$ has fewer than $(n+1-t)$ vertical arcs and is zero in $\mathbb{I}_{n-1}^{n+1-t} \mathbf{u}_{n-1, t-2}$. Therefore φ_3 is an isomorphism of \mathbb{A}_{n-1} -left

modules. In view of $\text{ind}_{S_{n-t}}^{S_{n+1-t}}(S^\lambda) \cong \bigoplus_{\lambda \sqsubset \beta} S^\beta$ we derive

$$\begin{aligned} \text{res}_{n-1}(\mathcal{M}_{\mathbb{A}_n}(\lambda)/F_n(\lambda)) &\cong \mathbb{I}_{n-1}^{n+1-t} \mathbf{u}_{n-1,t-2} \otimes_{S_{n+1-t}} \text{ind}_{S_{n-t}}^{S_{n+1-t}}(S^\lambda) \\ &\cong \bigoplus_{\lambda \sqsubset \beta} (\mathbb{I}_{n-1}^{n+1-t} \mathbf{u}_{n-1,t-2} \otimes_{S_{n+1-t}} S^\beta) \\ &\cong \bigoplus_{\lambda \sqsubset \beta} \mathcal{M}_{\mathbb{A}_{n-1}}(\beta) \end{aligned}$$

and the proof of the theorem is complete. \square

For \mathbb{L}_n , there exists no nontrivial space $G_n(\lambda)$ and Theorem 1 accordingly implies

Corollary 1. *Let $n, t \in \mathbb{N}$ and $\lambda \vdash (n-t)$ where $1 \leq t \leq n$. Then we have the isomorphism of \mathbb{L}_{n-1} -modules*

$$(3.15) \quad \bigoplus_{\alpha \sqsubseteq \lambda} \mathcal{M}_{\mathbb{L}_{n-1}}(\alpha) \cong \text{res}_{n-1}(\mathcal{M}_{\mathbb{L}_n}(\lambda)).$$

We proceed by studying induction in \mathbb{A}_n . Let us begin by remarking that the arguments of the following proof can easily be put into context with the localization and globalization functors [10, 4]. Since the latter are compatible with the quasi-hereditary structure of \mathbb{A}_n , in case of $x \neq 0$ one can obtain a more structural point of view.

Theorem 2. *Let $n, t \in \mathbb{N}$ and $\lambda \vdash (n-t)$ where $1 \leq t \leq n$. Then we have*

$$(3.16) \quad \text{ind}_{n+1}(\mathcal{M}_{\mathbb{A}_n}(\lambda)) \cong \text{res}_{n+1}(\mathcal{M}_{\mathbb{A}_{n+2}}(\lambda)).$$

Furthermore there exists the exact sequence of \mathbb{A}_{n+1} -modules

$$(3.17) \quad 0 \longrightarrow \bigoplus_{\alpha \sqsubseteq \lambda} \mathcal{M}_{\mathbb{A}_{n+1}}(\alpha) \longrightarrow \text{ind}_{n+1}(\mathcal{M}_{\mathbb{A}_n}(\lambda)) \longrightarrow \bigoplus_{\lambda \sqsubset \beta} \mathcal{M}_{\mathbb{A}_{n+1}}(\beta) \longrightarrow 0.$$

Proof. We first prove eq. (3.16). Suppose we have $\mathbf{a} \in \mathcal{A}_{n+2}$, with the property that its bottom vertices $(n+1)'$ and $(n+2)'$ are connected by a horizontal arc. Let $f_4(\mathbf{a})$ be the diagram obtained from \mathbf{a} by removing its bottom vertices $(n+1)'$, $(n+2)'$ together with their horizontal arc and moving its top-vertex $(n+2)$ to the bottom at position $(n+1)'$. It is straightforward to prove that for any $\mathcal{M}_{\mathbb{A}_n}(\lambda)$ the mapping

$$(3.18) \quad \begin{aligned} \varphi_4: \text{res}_{n+1}(\mathbb{A}_{n+2} \mathbf{e}_{n+1} \otimes_{\mathbb{A}_n} \mathcal{M}_{\mathbb{A}_n}(\lambda)) &\longrightarrow \mathbb{A}_{n+1} \otimes_{\mathbb{A}_n} \mathcal{M}_{\mathbb{A}_n}(\lambda) \\ \mathbf{a} \otimes w &\longmapsto f_4(\mathbf{a}) \otimes w, \end{aligned}$$

is an isomorphism of \mathbb{A}_{n+1} -modules. We proceed by showing

$$(3.19) \quad \mathbb{A}_{n+2}\mathfrak{e}_{n+1} \otimes_{\mathbb{A}_n} \mathcal{M}_{\mathbb{A}_n}(\lambda) \cong \mathcal{M}_{\mathbb{A}_{n+2}}(\lambda).$$

The key to eq. (3.19) is to prove that

$$(3.20) \quad \mathbb{A}_{n+2}\mathfrak{e}_{n+1} \otimes_{\mathbb{A}_n} \mathbb{I}_n^{n-t}\mathfrak{u}_{n,t} \cong \mathbb{I}_{n+2}^{n-t}\mathfrak{u}_{n+2,t+2}$$

is an isomorphism of \mathbb{A}_{n+2} -left modules. For this purpose we consider a tensor $\mathfrak{a}\mathfrak{e}_{n+1} \otimes \mathfrak{b}\mathfrak{u}_{n,t}$, where $\mathfrak{a}\mathfrak{e}_{n+1} \in \mathbb{A}_{n+2}\mathfrak{e}_{n+1}$ and $\mathfrak{b} \in \mathcal{A}_n^{n-t}$. Let $\mathfrak{r} \in \mathcal{A}_n^{n-t}$ be obtained from \mathfrak{b} as follows: we set $\text{bot}(\mathfrak{r}) = \text{top}(\mathfrak{b})$, $\text{top}(\mathfrak{r}) = \text{top}(\mathfrak{u}_{n,t})$ and choose the vertical \mathfrak{r} -arcs and $m \in \mathbb{Z}$ such that

$$(3.21) \quad x^m \mathfrak{r} \mathfrak{b} \mathfrak{u}_{n,t} = \mathfrak{u}_{n,t}.$$

Since the product $\mathfrak{r}^*\mathfrak{r}$ generates exactly t inner components, we obtain using eq. (3.21)

$$x^{-t+m} \mathfrak{r}^* \mathfrak{r} \mathfrak{b} \mathfrak{u}_{n,t} = \mathfrak{b} \mathfrak{u}_{n,t}.$$

Using $\text{bot}(\mathfrak{r}^*) = \text{top}(\mathfrak{r}) = \text{top}(\mathfrak{u}_{n,t})$, we compute

$$\begin{aligned} \mathfrak{a}\mathfrak{e}_{n+1} \otimes \mathfrak{b} \mathfrak{u}_{n,t} &= \mathfrak{a}\mathfrak{e}_{n+1} \otimes x^{-t+m} \mathfrak{r}^* \mathfrak{r} \mathfrak{b} \mathfrak{u}_{n,t} \\ &= \mathfrak{a}\mathfrak{e}_{n+1} x^{-t} \mathfrak{r}^* \otimes x^m \mathfrak{r} \mathfrak{b} \mathfrak{u}_{n,t} \\ &= \mathfrak{a}\mathfrak{e}_{n+1} x^{-t} \mathfrak{r}^* \otimes \mathfrak{u}_{n,t} \\ &= x^{-t} \mathfrak{a} \mathfrak{r}^* \mathfrak{e}_{n+1} \otimes \mathfrak{u}_{n,t} \\ &= \mathfrak{a}' \mathfrak{u}_{n,t} \mathfrak{e}_{n+1} \otimes \mathfrak{u}_{n,t}. \end{aligned}$$

Employing the just derived normal form for tensors, we are now in position to make the isomorphism of \mathbb{A}_{n+2} -left modules of eq. (3.20) explicit

$$\begin{aligned} \varphi_5: \mathbb{A}_{n+2}\mathfrak{e}_{n+1} \otimes_{\mathbb{A}_n} \mathbb{I}_n^{n-t}\mathfrak{u}_{n,t} &\longrightarrow \mathbb{I}_{n+2}^{n-t}\mathfrak{u}_{n+2,t+2} \\ \mathfrak{a}' \mathfrak{u}_{n,t} \mathfrak{e}_{n+1} \otimes \mathfrak{u}_{n,t} &\longmapsto \mathfrak{a}' \mathfrak{u}_{n+2,t+2}. \end{aligned}$$

Standard tensor identities imply

$$\begin{aligned} \mathbb{A}_{n+2}\mathfrak{e}_{n+1} \otimes_{\mathbb{A}_n} \mathcal{M}_{\mathbb{A}_n}(\lambda) &\cong (\mathbb{A}_{n+2}\mathfrak{e}_{n+1} \otimes_{\mathbb{A}_n} \mathbb{I}_n^{n-t}\mathfrak{u}_{n,t}) \otimes_{S_{n-t}} S^\lambda \\ &\cong \mathbb{I}_{n+2}^{n-t}\mathfrak{u}_{n+2,t+2} \otimes_{S_{n-t}} S^\lambda \\ &\cong \mathcal{M}_{\mathbb{A}_{n+2}}(\lambda). \end{aligned}$$

Now Claim 3 follows immediately

$$\begin{aligned} \text{ind}_{n+1}(\mathcal{M}_{\mathbb{A}_n}(\lambda)) &= \mathbb{A}_{n+1} \otimes_{\mathbb{A}_n} \mathcal{M}_{\mathbb{A}_n}(\lambda) \\ &\cong \text{res}_{n+1}(\mathbb{A}_{n+2}\mathfrak{e}_{n+1} \otimes_{\mathbb{A}_n} \mathcal{M}_{\mathbb{A}_n}(\lambda)) \\ &\cong \text{res}_{n+1}(\mathcal{M}_{\mathbb{A}_{n+2}}(\lambda)). \end{aligned}$$

Accordingly, the exact sequence of eq. (3.17) is immediately implied by Theorem 1 and the proof of the theorem is complete. \square

Corollary 2. *Let $n, t \in \mathbb{N}$ and $\lambda \vdash (n - t)$ where $1 \leq t \leq n$. Then we have*

$$(3.22) \quad \text{ind}_{n+1}(\mathcal{M}_{\mathbb{L}_n}(\lambda)) \cong \text{res}_{n+1}(\mathcal{M}_{\mathbb{L}_{n+2}}(\lambda)) \quad \text{and} \quad \bigoplus_{\alpha \sqsubseteq \lambda} \mathcal{M}_{\mathbb{L}_{n+1}}(\alpha) \cong \text{ind}_{n+1}(\mathcal{M}_{\mathbb{L}_n}(\lambda)).$$

4. SEMISIMPLICITY

The semisimplicity of \mathbb{L}_n is an immediate consequence of Proposition 1 and Proposition 2.

Theorem 3. *Suppose $x \neq 0$, then \mathbb{L}_n is semisimple.*

Proof. We showed in Proposition 2, that if \mathbb{L}_n is not semisimple, then there exist two partitions μ, λ , where $|\mu| < |\lambda| \leq n$ and a nontrivial morphism of \mathbb{L}_n -modules $\mathcal{M}_{\mathbb{L}_n}(\lambda) \xrightarrow{\varphi_n} \mathcal{M}_{\mathbb{L}_n}(\mu)$. The uniqueness of $\mathcal{N}_{\mathbb{L}_n}(\mu)$ implies that $\varphi_n(\mathcal{M}_{\mathbb{L}_n}(\lambda)) \subset \mathcal{N}_{\mathbb{L}_n}(\mu)$.

Claim. For $x \neq 0$ we have $\mathcal{N}_{\mathbb{L}_n}(\mu) = 0$.

In case of $\mu \vdash n$ this follows immediately from the irreducibility of the lift of the Specht module S^λ . Suppose next $\mu \vdash (n - t) < n$. Let $\mathbf{a} \in \mathcal{J}_n^{n-t}$, where $\text{bot}(\mathbf{a}) = \text{bot}(\mathbf{u}_{n,t})$ and let $v \in S^\mu$. For any $\mathbf{a} \otimes v \in \mathcal{M}_{\mathbb{L}_n}(\mu)$, there exists some $\sigma_0 \in S_{n-t}$ and some t -tuple (j_1, j_2, \dots, j_t) , where $1 \leq j_1 < j_2 < \dots < j_t \leq n$ such that $\mathbf{a}_{(j_h)_{h=1}^t} = \mathbf{a}\sigma_0$ has noncrossing vertical arcs and has top-vertex loops at j_1, \dots, j_t . The \mathcal{J}_n^{n-t} -diagram, $\mathbf{a}_{(j_h)_{h=1}^t}$ has the property $\mathbf{a} \otimes v = \mathbf{a}_{(j_h)_{h=1}^t} \otimes \sigma_0^{-1}v$ and any $u \in \mathbb{I}_n^{n-t} \mathbf{u}_{n,t} \otimes_{S_{n-1}} S^\mu$ can be written as

$$u = \sum_{1 \leq j_1 < j_2 < \dots < j_t \leq n} \mathbf{a}_{(j_h)_{h=1}^t} \otimes w_{(j_h)_{h=1}^t}.$$

For $U_t = \sum_{1 \leq i_1 < i_2 < \dots < i_t \leq n} \mathbf{u}_{i_1} \cdots \mathbf{u}_{i_t} \in \mathbb{I}_n^{n-t}$ we immediately obtain $U_t \cdot \mathbf{a}_{(j_h)_{h=1}^t} = x^t \mathbf{a}_{(j_h)_{h=1}^t}$. Indeed, only if (i_1, i_2, \dots, i_t) matches the tuple $(j_h)_{h=1}^t$ the factor x^t via the t -inner components of the graph $G(\mathbf{u}_{i_j} \cdots \mathbf{u}_{i_t}, \mathbf{a}_{(j_h)_{h=1}^t})$ is produced. In all other cases there exists a loop which traces back to the bottom row of $G'(\mathbf{u}_{i_j} \cdots \mathbf{u}_{i_t}, \mathbf{a}_{(j_h)_{h=1}^t})$ resulting in a zero in \mathbb{I}_n^{n-t} . Therefore, for any $u \in \mathcal{M}_{\mathbb{L}_n}(\mu)$

$$(4.1) \quad U_t \cdot u = x^t u$$

holds. Since $U_t \in \mathbb{I}_n^{n-t}$, $x \neq 0$ implies $\mathcal{N}_{\mathbb{L}_n}(\mu) = \{w \in \mathcal{M}_{\mathbb{L}_n}(\mu) \mid \mathbb{I}_n^{n-t} w = 0\} = 0$ and the Claim is proved.

The uniqueness of $\mathcal{N}_{\mathbb{L}_n}(\mu)$ as a maximal $\mathcal{M}_{\mathbb{L}_n}(\mu)$ module implies that any nontrivial morphism φ_n has the property $\varphi_n(\mathcal{M}_{\mathbb{L}_n}(\lambda)) \subset \mathcal{N}_{\mathbb{L}_n}(\mu)$. Therefore we arrive at $\varphi_n(\mathcal{M}_{\mathbb{L}_n}(\lambda)) = 0$, i.e. there exists no nontrivial morphism $\varphi_n: \mathcal{M}_{\mathbb{L}_n}(\lambda) \rightarrow \mathcal{M}_{\mathbb{L}_n}(\mu)$, whence \mathbb{L}_n is semisimple. \square

We next consider the algebra \mathbb{A}_n . According to Proposition 2, if \mathbb{A}_n is not semisimple then there exists the exact sequence

$$(4.2) \quad 0 \longrightarrow \mathcal{N}_{\mathbb{A}_n}(\lambda) \longrightarrow \mathcal{M}_{\mathbb{A}_n}(\lambda) \xrightarrow{\varphi_n} \mathcal{M}_{\mathbb{A}_n}(\mu),$$

where μ, λ are two partitions, such that $\lambda \vdash (n - t_\lambda)$ and $\mu \vdash (n - t_\mu)$, $t_\lambda < t_\mu$. In the next lemma we show that we can, without loss of generality, assume that $\lambda \vdash n$. Since $\mathcal{M}_{\mathbb{A}_n}(\lambda) \cong S^\lambda$ is irreducible this implies that we have an embedding $\varphi_n: S^\lambda \rightarrow \mathcal{M}_{\mathbb{A}_n}(\mu)$.

Lemma 2. *Suppose $x \neq 0$ and \mathbb{A}_n is not semisimple. Then there exists $n_1 \leq n$, two partitions $\lambda_1 \vdash n_1$, $\mu_1 \vdash n_1 - t_1$ and the short exact sequence*

$$(4.3) \quad 0 \longrightarrow S^{\lambda_1} \xrightarrow{\varphi_{n_1}} \mathcal{M}_{\mathbb{A}_{n_1}}(\mu_1).$$

Proof. If \mathbb{A}_n is not semisimple, then there exists $\lambda \vdash (n - t_\lambda)$, $\mu \vdash (n - t_\mu)$, where $t_\lambda < t_\mu$ and the exact sequence of eq. (4.2). Without loss of generality we may assume $0 < t_\lambda$. Theorem 1 guarantees the existence of the embedding $e_\lambda: \mathcal{M}_{\mathbb{A}_{n-1}}(\lambda) \rightarrow \mathcal{M}_{\mathbb{A}_n}(\lambda)$ and $e_\mu: \mathcal{M}_{\mathbb{A}_{n-1}}(\mu) \rightarrow \mathcal{M}_{\mathbb{A}_n}(\mu)$ given by $e_\lambda(\mathbf{a} \otimes v) = \mathbf{a}u_n \otimes v$ and $e_\mu(\mathbf{a} \otimes w) = \mathbf{a}u_n \otimes w$, respectively. We shall show that $\varphi_n: \mathcal{M}_{\mathbb{A}_n}(\lambda) \rightarrow \mathcal{M}_{\mathbb{A}_n}(\mu)$ induces a nontrivial morphism of \mathbb{A}_{n-1} -left modules via

$$(4.4) \quad \begin{array}{ccc} \mathcal{M}_{\mathbb{A}_n}(\lambda) & \xrightarrow{\varphi_n} & \mathcal{M}_{\mathbb{A}_n}(\mu) \\ \uparrow e_\lambda & & \uparrow e_\mu \\ \mathcal{M}_{\mathbb{A}_{n-1}}(\lambda) & \xrightarrow{\varphi_{n-1}} & \mathcal{M}_{\mathbb{A}_{n-1}}(\mu) \end{array}$$

Let $\mathbf{a} \otimes v \in \mathcal{M}_{\mathbb{A}_{n-1}}(\lambda)$, where $\mathbf{a} \in \mathcal{J}_{n-1}^{n-t_\lambda}$, $\text{bot}(\mathbf{a}) = \text{bot}(u_{n-1, t_\lambda-1})$ and $v \in S^\lambda$. Since φ_n is a morphism of \mathbb{A}_n -left modules we have $\varphi_n(u_n \cdot e_\lambda(\mathbf{a} \otimes v)) = u_n \cdot \varphi_n(e_\lambda(\mathbf{a} \otimes v))$ and in view of $\varphi_n(u_n \cdot e_\lambda(\mathbf{a} \otimes v)) = x \varphi_n(e_\lambda(\mathbf{a} \otimes v))$ we derive

$$(4.5) \quad u_n \varphi_n(e_\lambda(\mathbf{a} \otimes v)) = x \varphi_n(e_\lambda(\mathbf{a} \otimes v)).$$

We represent $\varphi_n(e_\lambda(\mathbf{a} \otimes v)) = \sum_r \mathbf{a}_r \otimes v_r$, where the \mathbf{a}_r are distinct $\mathcal{J}_n^{n-t_\mu}$ -diagrams, having noncrossing verticals, with $\text{bot}(\mathbf{a}_r) = \text{bot}(u_{n, t_\mu})$ and $v_r \in S^\mu$. Then we obtain

$$(4.6) \quad x^{-1} u_n \varphi_n(e_\lambda(\mathbf{a} \otimes v)) = x^{-1} \sum_r (u_n \mathbf{a}_r) \otimes v_r = \sum_r \mathbf{a}_r \otimes v_r.$$

Since different $\mathcal{J}_n^{n-t_\mu}$ -diagrams are by construction linear independent we can conclude from eq. (4.6), that each \mathbf{a}_r has a loop at top-vertex n . Therefore there exists for each $\varphi_n(e_\lambda(\mathbf{a} \otimes v)) = \sum_r \mathbf{a}_r \otimes v_r$, a unique element $\sum_r \mathbf{a}_r^\dagger \otimes v_r \in \mathcal{M}_{\mathbb{A}_{n-1}}(\mu)$, obtained by removing the vertices n and n' and their corresponding loops from each \mathbf{a}_r . Since $\mathcal{M}_{\mathbb{A}_{n-1}}(\lambda)$ is generated by tensors of the form $\mathbf{a} \otimes v$, φ_n induces the mapping

$$(4.7) \quad \begin{aligned} \varphi_{n-1}: \mathcal{M}_{\mathbb{A}_{n-1}}(\lambda) &\longrightarrow \mathcal{M}_{\mathbb{A}_{n-1}}(\mu) \\ \mathbf{a} \otimes v &\longmapsto \sum_r \mathbf{a}_r^\dagger \otimes v_j, \end{aligned}$$

with the property $e_\mu \cdot \varphi_{n-1} = \varphi_n \cdot e_\lambda$, i.e. φ_{n-1} makes the diagram in eq. (4.4) commutative. By construction, φ_{n-1} is a morphism of \mathbb{A}_{n-1} -left modules.

Claim. We have $w \in \mathcal{N}_{\mathbb{A}_{n-1}}(\lambda)$ if and only if $e_\lambda(w) \in \mathcal{N}_{\mathbb{A}_n}(\lambda)$.

Suppose first $w = \sum_i \mathbf{a}_i \otimes v_i \notin \mathcal{N}_{\mathbb{A}_{n-1}}(\lambda)$. According to eq. (2.5), there exists some $\mathbf{b}_0 \in \mathcal{J}_{n-1}^{n-t_\lambda}$ such that

$$\mathbf{b}_0 \cdot \sum_i \mathbf{a}_i \otimes v_i = \sum_i (\mathbf{b}_0 \mathbf{a}_i) \otimes v_i = \tilde{\mathbf{b}}_0 \otimes \sum_i \delta_{\mathbf{b}_0 \mathbf{a}_i} x^{\ell(\mathbf{b}_0, \mathbf{a}_i)} \sigma_{\mathbf{b}_0, \mathbf{a}_i} v_i \neq 0.$$

This equation implies in the \mathbb{A}_n -module $\mathcal{M}_{\mathbb{A}_n}(\lambda)$

$$\mathbf{b}_0 \cdot \sum_i \mathbf{a}_i \mathbf{u}_n \otimes v_i = \sum_i (\mathbf{b}_0 \mathbf{u}_n \mathbf{a}_i) \otimes v_i$$

where $\mathbf{b}_0 \mathbf{u}_n \in \mathcal{J}_n^{n-t_\lambda}$. In view of $\ell(\mathbf{b}_0 \mathbf{u}_n, \mathbf{a}_i) = \ell(\mathbf{b}_0, \mathbf{a}_i)$ and $\mathbf{b}_0 \mathbf{a}_i = x^{\ell(\mathbf{b}_0, \mathbf{a}_i)} \tilde{\mathbf{b}}_0 \sigma_{\mathbf{b}_0, \mathbf{a}_i}$, where $\sigma_{\mathbf{b}_0, \mathbf{a}_i} \in S_{n-t_\lambda}$ we obtain

$$\mathbf{b}_0 \cdot \sum_i \mathbf{a}_i \mathbf{u}_n \otimes v_i = \tilde{\mathbf{b}}_0 \mathbf{u}_n \otimes \sum_i \delta_{\mathbf{b}_0 \mathbf{a}_i} x^{\ell(\mathbf{b}_0, \mathbf{a}_i)} \sigma_{\mathbf{b}_0, \mathbf{a}_i} v_i \neq 0.$$

I.e. we have shown $w \notin \mathcal{N}_{\mathbb{A}_{n-1}}(\lambda) \implies e_\lambda(w) \notin \mathcal{N}_{\mathbb{A}_n}(\lambda)$. Second suppose $e_\lambda(w) \notin \mathcal{N}_{\mathbb{A}_n}(\lambda)$. Then there exists some $\mathbf{b}_0 \in \mathcal{J}_n^{n-t_\lambda}$ such that

$$\mathbf{b}_0 \cdot \sum_i \mathbf{a}_i \mathbf{u}_n \otimes v_i = \sum_i (\mathbf{b}_0 \mathbf{u}_n) \mathbf{a}_i \otimes v_i \neq 0$$

and $\mathbf{b}_0 \mathbf{u}_n$ is the scalar multiple of a diagram $\mathbf{r} \in \mathcal{J}_n^{n-t_\lambda}$, having $\text{top}(\mathbf{r}) = \text{top}(\mathbf{b}_0)$ and a loop at n' . We accordingly compute

$$\mathbf{b}_0 \cdot \sum_i \mathbf{a}_i \mathbf{u}_n \otimes v_i = x^s \sum_i \mathbf{r} \mathbf{a}_i \otimes v_i = x^s \tilde{\mathbf{r}} \otimes \sum_i \delta_{\mathbf{r} \mathbf{a}_i} x^{\ell(\mathbf{r}, \mathbf{a}_i)} \sigma_{\mathbf{r}, \mathbf{a}_i} v_i \neq 0,$$

where $\tilde{\mathbf{r}}$ is given by $\mathbf{r} \mathbf{a}_i = x^{\ell(\mathbf{r}, \mathbf{a}_i)} \delta_{\mathbf{r} \mathbf{a}_i} \tilde{\mathbf{r}} \sigma_{\mathbf{r}, \mathbf{a}_i}$. We may assume that \mathbf{r} has a loop at top-vertex n , since this feature does not affect the term $w_1 = \sum_i \delta_{\mathbf{r} \mathbf{a}_i} x^{\ell(\mathbf{r}, \mathbf{a}_i)} \sigma_{\mathbf{r}, \mathbf{a}_i} v_i$. By construction, $\tilde{\mathbf{r}}$ has then also a

loop at n and there exists a $\mathbf{c} \in \mathcal{J}_{n-1}^{n-t_\lambda}$ with the property $\mathbf{r} = \mathbf{c}u_n$ in $\mathcal{J}_n^{n-t_\lambda}$. In view of $\delta_{\mathbf{r}\mathbf{a}_i} = \delta_{\mathbf{c}\mathbf{a}_i}$, $\sigma_{\mathbf{r},\mathbf{a}_i} = \sigma_{\mathbf{c},\mathbf{a}_i}$ and $\ell(\mathbf{r}, \mathbf{a}_i) = \ell(\mathbf{c}, \mathbf{a}_i)$ we obtain

$$\mathbf{c} \cdot \sum_i \mathbf{a}_i \otimes v_i = \sum_i (\mathbf{c}\mathbf{a}_i) \otimes v_i = \tilde{\mathbf{c}} \otimes \sum_i \delta_{\mathbf{c}\mathbf{a}_i} x^{\ell(\mathbf{c},\mathbf{a}_i)} \sigma_{\mathbf{c},\mathbf{a}_i} v_i = \tilde{\mathbf{c}} \otimes w_1 \neq 0.$$

That is, we have proved $e_\lambda(w) \notin \mathcal{N}_{\mathbb{A}_n}(\lambda) \implies w \notin \mathcal{N}_{\mathbb{A}_{n-1}}(\lambda)$ and the Claim follows.

Using $e_\mu \cdot \varphi_{n-1} = \varphi_n \cdot e_\lambda$, we can now immediately conclude $\ker(\varphi_{n-1}) = \mathcal{N}_{\mathbb{A}_{n-1}}(\lambda)$. Indeed, if $w \in \ker(\varphi_{n-1})$ then $e_\lambda(w) \in \ker(\varphi_n) = \mathcal{N}_{\mathbb{A}_n}(\lambda)$, whence $w \in \mathcal{N}_{\mathbb{A}_{n-1}}(\lambda)$. If $w \in \mathcal{N}_{\mathbb{A}_{n-1}}(\lambda)$, then $e_\lambda(w) \in \mathcal{N}_{\mathbb{A}_n}(\lambda)$, whence $e_\mu \cdot \varphi_{n-1}(w) = 0$, from which $\varphi_{n-1}(w) = 0$, i.e. $w \in \ker(\varphi_{n-1})$ follows. Therefore we have the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{N}_{\mathbb{A}_n}(\lambda) & \longrightarrow & \mathcal{M}_{\mathbb{A}_n}(\lambda) & \xrightarrow{\varphi_n} & \mathcal{M}_{\mathbb{A}_n}(\mu) \\ & & \uparrow e_\lambda & & \uparrow e_\lambda & & \uparrow e_\mu \\ 0 & \longrightarrow & \mathcal{N}_{\mathbb{A}_{n-1}}(\lambda) & \longrightarrow & \mathcal{M}_{\mathbb{A}_{n-1}}(\lambda) & \xrightarrow{\varphi_{n-1}} & \mathcal{M}_{\mathbb{A}_{n-1}}(\mu). \end{array}$$

Iterating the above construction we arrive, after t_λ steps, at some $\lambda_1 \vdash n_1$, $\mu_1 \vdash (n - t_1)$ and the exact sequence $0 \longrightarrow \mathcal{N}_{\mathbb{A}_{n_1}}(\lambda_1) \longrightarrow \mathcal{M}_{\mathbb{A}_{n_1}}(\lambda_1) \longrightarrow \mathcal{M}_{\mathbb{A}_{n_1}}(\mu_1)$. Since $\lambda_1 \vdash n_1$ we have $\mathcal{N}_{\mathbb{A}_{n_1}}(\lambda_1) = 0$ and $\mathcal{M}_{\mathbb{A}_{n_1}}(\lambda_1) \cong S^{\lambda_1}$, whence Lemma 2. \square

Our next result establishes further restrictions on the embedding $0 \longrightarrow S^\lambda \longrightarrow \mathcal{M}_{\mathbb{A}_n}(\mu)$ in terms of the partition μ . Using the exact sequence for the restriction functor of Theorem 1 we shall prove the existence of such an embedding with the additional property $\mu_2 \vdash (n-1)$ or $\mu_2 \vdash (n-2)$. The result is in analogy to the Brauer algebra case proved in [4].

Lemma 3. *Suppose $\varphi_{n_1}: S^{\lambda_1} \longrightarrow \mathcal{M}_{\mathbb{A}_{n_1}}(\mu_1)$ is an embedding where $\lambda_1 \vdash n_1$ and $\mu_1 \vdash n_1 - t_1$. Then for \mathbb{A}_n there exist $n_2 \leq n_1$, a pair of partitions (λ_2, μ_2) and an embedding $S^{\lambda_2} \longrightarrow \mathcal{M}_{\mathbb{A}_{n_2}}(\mu_2)$, such that $\lambda_2 \vdash n_2$, $\mu_2 \vdash (n_2 - 1)$ or $\mu_2 \vdash (n_2 - 2)$.*

Proof. Since $\text{res}_{S_{n_1-1}}(S^{\lambda_1}) \cong \bigoplus_{\nu \sqsubset \lambda_1} S^\nu$ we obtain for some $\nu \sqsubset \lambda_1$, $\nu \vdash (n_1 - 1)$, the embedding $\varphi_\nu: S^\nu \longrightarrow \text{res}_{n_1-1}(\mathcal{M}_{\mathbb{A}_{n_1}}(\mu_1))$. An interpretation of $\text{res}_{n_1-1}(\mathcal{M}_{\mathbb{A}_{n_1}}(\mu_1))$ is given via Theorem 1 in terms of the exact sequence

$$(4.8) \quad 0 \longrightarrow \bigoplus_{\alpha_1 \sqsubseteq \mu_1} \mathcal{M}_{n_1-1}(\alpha_1) \longrightarrow \text{res}_{n_1-1}(\mathcal{M}_{\mathbb{A}_{n_1}}(\mu_1)) \longrightarrow \bigoplus_{\mu_1 \sqsubset \beta_1} \mathcal{M}_{n_1-1}(\beta_1) \longrightarrow 0.$$

Suppose we have $\varphi_\nu(S^\nu) \subset F_{n_1}(\mu_1) \cong \bigoplus_{\alpha_1 \sqsubseteq \mu_1} \mathcal{M}_{n_1-1}(\alpha_1)$. Then the irreducibility of S^ν implies the embedding $S^\nu \longrightarrow \mathcal{M}_{n_1-1}(\alpha_1)$ for some $\alpha_1 \sqsubseteq \mu_1$. Otherwise, we have $\varphi_\nu(S^\nu) \not\subset F_{n_1}(\mu_1)$. The

irreducibility of $\varphi_\nu(S^\nu)$ guarantees

$$(\varphi_\nu(S^\nu) \oplus F_{n_1}(\mu_1)) / F_{n_1}(\mu_1) \cong \varphi_\nu(S^\nu).$$

In view of eq. (4.8) we have

$$\text{res}_{n_1-1}(\mathcal{M}_{\mathbb{A}_{n_1}}(\mu_1)) / F_{n_1}(\mu_1) \cong \bigoplus_{\mu_1 \sqsubset \beta_1} \mathcal{M}_{n_1-1}(\beta_1)$$

which implies an embedding $S^\nu \longrightarrow \mathcal{M}_{n_1-1}(\beta_1)$, for some $\mu_1 \sqsubset \beta_1$.

Therefore we have the following situation: each iteration of the above argument reduces the size of the partition $\lambda_1 \vdash n_1$ by one and an analogous reduction of the partition μ_1 can occur at most $(n_1 - t_1) < n_1$ times. Any further iteration cannot decrease the size of μ_1 , while decreasing the size of λ_1 . That is, iteration produces a pair (λ_2, μ_2) where $\lambda_2 \vdash n_2$ and $\mu_2 \vdash (n_2 - h)$, where $h = 1$ or $h = 2$. Indeed, for $h = 2$, i.e. $\mu_2 \vdash n_2 - 2$, further reduction can generate the trivial embedding $S^\nu \longrightarrow S^\nu$, i.e. we derive, using the above notation, $\nu = \beta_1$, for $\nu \sqsubset \lambda_2$. Therefore further reduction is in general not possible and we have shown that iteration of the above process leads to a pair of partitions (λ_2, μ_2) with the properties $\lambda_2 \vdash n_2$ and $\mu_2 \vdash (n_2 - 1)$ or $\mu_2 \vdash (n_2 - 2)$. \square

Now we are in position to prove our main result:

Theorem 4. *Suppose $x \neq 0$. If $x \notin \mathbb{Z}$, then the algebra \mathbb{A}_n is semisimple.*

Proof. According to Proposition 2, if \mathbb{A}_n is not semisimple there exists a nontrivial morphism $\varphi_n: \mathcal{M}_{\mathbb{A}_n}(\lambda) \longrightarrow \mathcal{M}_{\mathbb{A}_n}(\mu)$ with $\ker(\varphi_n) = \mathcal{N}_{\mathbb{A}_n}(\lambda)$.

In view of Lemma 2 and Lemma 3 we can, without loss of generality, assume that there exists an embedding $\varphi_n: S^\lambda \longrightarrow \mathcal{M}_{\mathbb{A}_n}(\mu)$, where $\lambda \vdash n$ and either $\mu \vdash (n - 1)$ or $\mu \vdash (n - 2)$. According to Proposition 1, $\mathcal{N}_{\mathbb{A}_n}(\mu)$ is the unique maximal $\mathcal{M}_{\mathbb{A}_n}(\mu)$ -submodule. Therefore $\varphi_n(S^\lambda) \subset \mathcal{N}_{\mathbb{A}_n}(\mu)$, i.e. we have the embedding $\varphi_n: S^\lambda \longrightarrow \mathcal{N}_{\mathbb{A}_n}(\mu)$. In the following we distinguish the two cases $\mu \vdash (n - 1)$ and $\mu \vdash (n - 2)$.

Case 1: $\mu \vdash (n - 1)$. We prove that $x \neq 0$ implies $\mathcal{N}_{\mathbb{A}_n}(\mu) = 0$. Let $\mathbf{a} \in \mathcal{J}_n^{n-1}$, where $\text{bot}(\mathbf{a}) = \text{bot}(\mathbf{u}_{n,1})$ and let $v \in S^\mu$. For any $\mathbf{a} \otimes v \in \mathcal{M}_{\mathbb{A}_n}(\mu)$, there exists some $\sigma_0 \in S_{n-1}$ and some index $1 \leq j \leq n$ such that $\mathbf{a}_j = \mathbf{a}\sigma_0$ has noncrossing vertical arcs and has its unique, top-vertex loop at j . \mathbf{a}_j has the property $\mathbf{a} \otimes v = \mathbf{a}_j \otimes \sigma_0^{-1}v$ and any $u \in \mathbb{I}_n^{n-1}\mathbf{u}_{n,1} \otimes_{S_{n-1}} S^\mu$ can be written as $u = \sum_j \mathbf{a}_j \otimes w_j$. Let $U_1 = \sum_i \mathbf{u}_i$. Then $U_1 \in \mathbb{I}_n^{n-1}$ and any \mathbf{a}_j satisfies the eigenvector equation $U_1 \cdot \mathbf{a}_j = x \mathbf{a}_j$. Let $u = \sum_j \mathbf{a}_j \otimes w_j \in \mathcal{N}_{\mathbb{A}_n}(\mu)$. Since $U_1 \in \mathbb{I}_n^{n-1}$, the action of U_1 on $\mathcal{N}_{\mathbb{A}_n}(\mu)$ is

trivial, i.e.


$$(4.9) \quad U_1 \cdot u = \sum_j (U_1 \cdot \mathbf{a}_j) \otimes w_j = \sum_i x \mathbf{a}_j \otimes w_j = x u = 0,$$

which implies, in view of $x \neq 0$, $\mathcal{N}_{\mathbb{A}_n}(\mu) = 0$.

Case 2: $\mu \vdash (n-2)$. For each diagram $\mathbf{a} \in \mathcal{J}_n^{n-2}$, such that $\text{bot}(\mathbf{a}) = \text{bot}(\mathbf{u}_{n,2})$ there exist a pair of indices, $i < j$ and a permutation $\sigma_0 \in S_{n-2}$ such that either $\mathbf{a}\sigma_0 = \mathbf{a}_{i,j}^\cap$ or $\mathbf{a}\sigma_0 = \mathbf{a}_{i,j}^\circ$ holds. Here $\mathbf{a}_{i,j}^\cap \in \mathcal{J}_n^{n-2}$ has noncrossing verticals, a horizontal arc connecting i and j and $\text{bot}(\mathbf{a}_{i,j}^\cap) = \text{bot}(\mathbf{u}_{n,2})$. Analogously, $\mathbf{a}_{i,j}^\circ \in \mathcal{J}_n^{n-2}$ has noncrossing verticals, two loops at i, j and $\text{bot}(\mathbf{u}_{n,2})$. We can write each tensor $\mathbf{a} \otimes w$, where $\mathbf{a} \in \mathcal{J}_n^{n-2}$ with $\text{bot}(\mathbf{a}) = \text{bot}(\mathbf{u}_{n,2})$ and $w \in S^\mu$, uniquely as either $\mathbf{a}_{i,j}^\cap \otimes \sigma_0^{-1}w$ or $\mathbf{a}_{i,j}^\circ \otimes \sigma_0^{-1}w$. Let $g: \mathcal{M}_{\mathbb{A}_n}(\mu) \rightarrow \mathcal{M}_{\mathbb{A}_n}(\mu)$ be the involution given via linear extension of $g(\mathbf{a}_{i,j}^\circ \otimes w) = \mathbf{a}_{i,j}^\cap \otimes w$ and $g(\mathbf{a}_{i,j}^\cap \otimes w) = \mathbf{a}_{i,j}^\circ \otimes w$. Furthermore, let $\mathbf{v}_{i,j} \in \mathcal{J}_n^{n-2}$ be the diagram having straight verticals except of a horizontal arc connecting the top-vertices i, j and two loops at the bottom vertices i', j' , respectively. We introduce

$$(4.10) \quad U_2 = \sum_{i < j} \mathbf{u}_i \mathbf{u}_j, \quad V_2 = \sum_{i < j} \mathbf{v}_{i,j} \quad \text{and} \quad H_2 = \sum_{i < j} \mathbf{h}_{i,j},$$

where $\mathbf{h}_{i,j} \in \mathcal{J}_n^{n-2}$ has straight vertical arcs except of the top-vertices i, j and bottom-vertices i', j' , which are connected by a horizontal arc, respectively. We observe $U_2, V_2, H_2 \in \mathbb{I}_n^{n-2}$.



As for the action of U_2 , a routine computation yields $U_2 \cdot \mathbf{a}_{i,j}^\cap = x \mathbf{a}_{i,j}^\circ$ and $U_2 \cdot \mathbf{a}_{i,j}^\circ = x^2 \mathbf{a}_{i,j}^\cap$. Similarly we obtain for V_2 , $V_2 \cdot \mathbf{a}_{i,j}^\cap = x \mathbf{a}_{i,j}^\cap$ and $V_2 \cdot \mathbf{a}_{i,j}^\circ = x^2 \mathbf{a}_{i,j}^\cap$. Let $\tau_{(i,j)}$ act on the diagram $\mathbf{a}_{i,j}^\cap$ as the transposition $(i, j) \in S_n$ from the left and $\tilde{\tau}_{(a,b)}$ as transposition $(a, b) \in S_{n-2}$, from the right, respectively. Then

$$(4.11) \quad H_2 \cdot \mathbf{a}_{i,j}^\circ = x \mathbf{a}_{i,j}^\cap$$

$$(4.12) \quad H_2 \cdot \mathbf{a}_{i,j}^\cap = \left((x-1) + \sum_{i < j} \tau_{(i,j)} - \sum_{a < b} \tilde{\tau}_{(a,b)} \right) \mathbf{a}_{i,j}^\cap,$$

where eq. (4.12) holds according to [4], Lemma 2, p.655. We write an element $v \in \mathcal{N}_{\mathbb{A}_n}(\mu)$ as

$$v = \sum_{i,j} \mathbf{a}_{i,j}^\cap \otimes r_{i,j} + \sum_{i,j} \mathbf{a}_{i,j}^\circ \otimes s_{i,j}$$

and set $v^\cap = \sum_{i,j} \mathfrak{a}_{i,j}^\cap \otimes r_{i,j}$ and $v^\circ = \sum_{i,j} \mathfrak{a}_{i,j}^\circ \otimes s_{i,j}$. Since $(H_2 - x^{-1}V_2) \in \mathbb{I}_n^{n-2}$, we obtain

$$\begin{aligned} (H_2 - x^{-1}V_2) \cdot (v^\cap + v^\circ) &= H_2 \cdot v^\cap + H_2 \cdot v^\circ - x^{-1}V_2 \cdot v^\cap - x^{-1}V_2 \cdot v^\circ \\ &= H_2 \cdot v^\cap + xg(v^\circ) - v^\cap - xg(v^\circ) \\ &= H_2 \cdot v^\cap - v^\cap. \end{aligned}$$

Suppose now there exists some $0 \neq v_0 \in \varphi_n(S^\lambda) \subset \mathcal{N}_{\mathbb{A}_n}(\mu)$ such that $v_0^\cap \neq 0$ and $v_0^\circ \neq 0$. Since $\varphi_n(S^\lambda)$ is an irreducible S_n -module and the S_n -action cannot change a horizontal arc into a pair of loops, for any $0 \neq v \in \varphi_n(S^\lambda) \subset \mathcal{N}_{\mathbb{A}_n}(\mu)$, $v^\cap \neq 0$ and $v^\circ \neq 0$ holds. Therefore if there exists some $0 \neq v_0 \in \varphi_n(S^\lambda) \subset \mathcal{N}_{\mathbb{A}_n}(\mu)$ such that $v_0^\cap \neq 0$ and $v_0^\circ \neq 0$, then we have for any $0 \neq v \in \varphi_n(S^\lambda)$, $(H_2 - 1) \cdot v^\cap = 0$, i.e.

$$(4.13) \quad \left((x-1) + \sum_{i < j} \tau_{(i,j)} - \sum_{a < b} \tilde{\tau}_{(a,b)} - 1 \right) \cdot v^\cap = 0.$$

We proceed by studying the action of $\sum_{i < j} \tau_{(i,j)}$ and $\sum_{a < b} \tilde{\tau}_{(a,b)}$ on the set

$$(4.14) \quad \varphi_n^\cap(S^\lambda) = \{v^\cap \mid v^\cap + v^\circ \in \varphi(S^\lambda)\}.$$

The \mathbb{A}_n -module $\mathcal{M}_{\mathbb{A}_n}(\mu)$ can be regarded as a $S_n \times S_{n-2}$ -left module via

$$(4.15) \quad (\sigma, \sigma') \cdot (\mathfrak{a} \otimes w) = \sigma \cdot (\mathfrak{a} \otimes \sigma' w)$$

and $\sigma \cdot (\mathfrak{a} \otimes \sigma' w) = (\sigma \mathfrak{a} \sigma') \otimes w$ shows that the action of eq. (4.12) and eq. (4.15) coincide. Furthermore, $\varphi_n(S^\lambda)$ becomes via eq. (4.15) a $S_n \times S_{n-2}$ -submodule of $\mathcal{M}_{\mathbb{A}_n}(\mu)$ and induces an $S_n \times S_{n-2}$ action on the set $\varphi_n^\cap(S^\lambda)$ via $(\sigma, \sigma') \cdot (\mathfrak{a}_{i,j}^\cap \otimes w) = \sigma \cdot (\mathfrak{a}_{i,j}^\cap \otimes \sigma' w)$. Accordingly, $\varphi_n^\cap(S^\lambda)$ can be considered as a $S_n \times S_{n-2}$ -module and the projection

$$(4.16) \quad \pi_1 : \varphi_n(S^\lambda) \longrightarrow \varphi_n^\cap(S^\lambda), \quad (v^\cap + v^\circ) \mapsto v^\cap,$$

establishes an isomorphism of $S_n \times S_{n-2}$ -modules. Indeed, only injectivity needs to be proved. Using $x \neq 0$, $U_2 \in \mathbb{I}_n^{n-2}$ and $(v^\cap + v^\circ) \in \mathcal{N}_{\mathbb{A}_n}(\lambda)$, injectivity follows from

$$x^{-1}U_2 \cdot (v^\cap + v^\circ) = g(v^\cap) + xv^\circ = 0,$$

i.e. $v^\circ = -x^{-1}g(v^\cap)$. Obviously, $\sum_{i < j} \tau_{(i,j)}$ and $\sum_{a < b} \tilde{\tau}_{(a,b)}$ are contained in the centers of the group algebras $F[S_n]$ and $F[S_{n-2}]$, respectively and Schur's Lemma implies that they act as homotheties on irreducible representations. Since $\varphi_n(S^\lambda)$ embeds into the $S^\lambda \otimes S^\mu$ -component of $\mathcal{M}_{\mathbb{A}_n}(\mu)$, the particular values of $\sum_{i < j} \tau_{(i,j)}$ and $\sum_{a < b} \tilde{\tau}_{(a,b)}$ are given by [12]

$$(4.17) \quad \sum_{i < j} \tau_{(i,j)} = \sum_{p \in [\lambda]} c(p) \quad \text{and} \quad \sum_{a < b} \tilde{\tau}_{(a,b)} = \sum_{p \in [\mu]} c(p).$$

Since $\varphi_n(S^\lambda) \subset \mathcal{N}_{\mathbb{A}_n}(\mu)$ we obtain

$$(4.18) \quad \forall v^\cap \in \varphi_n^\cap(S^\lambda); \quad \left((x-1) + \sum_{p \in [\lambda]} c(p) - \sum_{p \in [\mu]} c(p) - 1 \right) v^\cap = 0,$$

which implies

$$(4.19) \quad (x-1) + \sum_{p \in [\lambda]} c(p) - \sum_{p \in [\mu]} c(p) - 1 = 0.$$

Since the content $c(p)$ is an integer, eq. (4.19) implies $x \in \mathbb{Z}$. It thus remains to consider the cases $v^\cap = 0$ or $v^\circ = 0$. The case of $v^\circ = 0$ is due to [4]. In analogy we derive, using the action of H_2 on $\varphi_n(S^\lambda)$

$$(4.20) \quad \forall v \in \varphi_n(S^\lambda); \quad H_2 \cdot v = \left((x-1) + \sum_{p \in [\lambda]} c(p) - \sum_{p \in [\mu]} c(p) \right) \cdot v = 0,$$

which implies $(x-1) + \sum_{p \in [\lambda]} c(p) - \sum_{p \in [\mu]} c(p) = 0$. This immediatly allows us to conclude $x \in \mathbb{Z}$. In case of $v^\cap = 0$ we obtain for any $v \in \varphi_n(S^\lambda)$

$$(4.21) \quad U_2 \cdot v = x^2 v = 0,$$

which is, in view of $x \neq 0$ impossible.

We have therefore showed that in case of $\mu \vdash (n-1)$, $x \neq 0$ implies $\mathcal{N}_{\mathbb{A}_n}(\mu) = 0$. Since $\mathcal{N}_{\mathbb{A}_n}(\mu)$ is the unique, maximal $\mathcal{M}_{\mathbb{A}_n}(\lambda)$ -submodule, there cannot exist an embedding $\varphi_n: S^\lambda \rightarrow \mathcal{M}_{\mathbb{A}_n}(\mu)$. In case of $\mu \vdash (n-2)$, our proof guarantees that for $x \notin \mathbb{Z}$, there exists no embedding $\varphi_n: S^\lambda \rightarrow \mathcal{M}_{\mathbb{A}_n}(\mu)$, whence \mathbb{A}_n is semisimple. \square

Acknowledgments. We are grateful to Andreas Dress and Jing Qin for helpful discussions. This work was supported by the 973 Project, the PCSIRT Project of the Ministry of Education, the Ministry of Science and Technology, and the National Science Foundation of China.

REFERENCES

- [1] A. Berele. A schensted-type correspondence for the symplectic group. *J. Comb. Th. Series A*, pages 320–328, 1986.
- [2] J. Birman and H. Wenzl. Braids, link polynomials and a new algebra. *Trans. Amer. Math. Soc.*, 313:249–273, 1989.
- [3] R. Brauer. On algebras which are connected with the semisimple continuous groups. *Math.*, 38:857–872, 1937.

- [4] W.F. Doran IV, D.B. Wales, and P.J. Hanlon. On the semisimplicity of the brauer centralizer algebras. *J. Algebra*, 211:647–685, 1999.
- [5] P. Hanlon and D. Wales. Eigenvalues connected with brauer’s centralizer algebras. *J. Algebra*, 121:446–476, 1989.
- [6] P. Hanlon and D. Wales. On the decomposition of brauer’s centralizer algebras. *J. Algebra*, 121:404–445, 1989.
- [7] P. Hanlon and D. Wales. Computing the discriminants of brauer’s centralizer algebras. *Math. Comp.*, 54:771–796, 1990.
- [8] Qin J. Jin E.Y and Reidys M.C. Combinatorics of rna structures with pseudoknots. *Bull.Math.Biol.*, 70:45–67, 2008.
- [9] V.F.R. Jones. Index for subfactors. *Inv. Math.*, 72:1–25, 1983.
- [10] P. Martin. The structure of the partition algebras. *J. Algebra*, 183:319–358, 1996.
- [11] H. Rui. A criterion on the semisimple brauer algebras. *J. Combin. Theory Ser. A*, 111:78–88, 2005.
- [12] B. Sagan. *The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions*. Springer, 1991.
- [13] N.E. Samra and R. King. Dimensions of irreducible representations of the classical lie groups. *J. Phys. A:Math. Gen.*, 12:2317–2328, 1979.
- [14] J.P. Serre. *Linear Representations of Finite Groups*. Springer, 1977.
- [15] S. Sundaram. On the combinatorics of representations of $sp(2n, \mathbb{C})$. *Trans. Amer. Math. Soc.*, 313:249–273, 1989.
- [16] H. Wenzl. On the structure of brauer’s centralizer algebras. *Ann. Math*, 128:173–193, 1988.

CENTER FOR COMBINATORICS, LPMC-TJKLC, NANKAI UNIVERSITY, TIANJIN 300071, P.R. CHINA, PHONE: *86-22-2350-6800, FAX: *86-22-2350-9272

E-mail address: reidys@nankai.edu.cn